



Original Article

Seasonal Decomposition of Sexual Victimization-Related Cases in Hyderabad, Pakistan

Aisha Rasheed^{1*}, Ummara Munir², Nadia Aslam¹, Abdul Samad¹, Muhammad Rafique Shaikh³ and Isirat Bibi¹¹Department of Forensic Medicine and Toxicology, Liaquat University of Medical and Health Sciences, Jamshoro, Pakistan²Department of Forensic Medicine and Toxicology, Sheikh Zayed Medical College, Rahim Yar Khan, Pakistan³Department of Forensic Medicine and Toxicology, Chandka Medical College, Shaheed Mohtarma Benazir Bhutto Medical University, Larkana, Pakistan

ARTICLE INFO

Keywords:

Sexual Assault, Seasonal Analysis, Casualties, Rape, Sodomy

How to Cite:

Rasheed, A., Munir, U., Aslam, N., Samad, A., Shaikh, M. R., & Bibi, I. (2024). Seasonal Decomposition of Sexual Victimization-Related Cases in Hyderabad, Pakistan: Sexual Victimization-Related Cases in Hyderabad. *Pakistan Journal of Health Sciences*, 5(04). <https://doi.org/10.54393/pjhs.v5i04.1392>

*Corresponding Author:

Aisha Rasheed

Department of Forensic Medicine and Toxicology, Liaquat University of Medical and Health Sciences, Jamshoro, Pakistan
aeshakhalid@yahoo.comReceived Date: 28th March, 2024Acceptance Date: 27th April, 2024Published Date: 30th April, 2024

ABSTRACT

The issue of sexual assault is a serious concern that contributes to abuses of human rights and risks to public health. Evaluating situations like these is absolutely necessary in order to make the judicial process easier for both the victim and the person who committed the crime. When the seasonality of sexual assault-related incidents is determined, it can provide the community and law enforcement authorities with information that can help them implement preventative measures. **Objective:** To assess the seasonality of sexual assault-related cases at casualty department of a teaching hospital in Hyderabad, Pakistan. **Methods:** In current descriptive/prospective study, ninety two sexual assault-related medico-legal cases (MLCs) were included. This study based on the seasonality of sexual assault-related MLCs using seasonal decomposition of three-year from January 2020 to December 2022 data from a leading tertiary care teaching hospital in the Hyderabad region. **Results:** Most of the cases (N=42) were reported in the year 2021. The percentage of sexual assault-related MLCs was found to be concentrated from June to August (10.86% to 14.13%). The highest number of sexual assault-related MLCs occurred in July. The seasonal decomposition also showed pronounced peaks in summer, with intense peaks in July for rape and total sexual assault cases. **Conclusions:** The number of sexual assault-related cases was aggregated in mid-year, showing a higher trend in summer season. Efficient preventive measures in certain seasons and education of vulnerable groups can minimize sexual violence in the community.

INTRODUCTION

Medical practice involves various aspects of human activities. Among them, one important part is related to questions of legal matters. The situations related to legal issues arise in multiple contexts including personal injury claims, medical malpractice lawsuits, criminal cases, and workers' compensation claims [1]. Sexual abuse represents a public health and human right problem that is growing in cultures of silence [2]. The reporting of sexual assault and related medico-legal cases (MLCs) is influenced by the norms and socio-cultural taboos of Pakistan [3]. The medical examination of sexual violence-related crime incidence is crucial in facilitating the justice

system for the victim and offender. Most forms of criminal activity are generally observed to display seasonal variations, although these patterns may vary among different types of crimes. Seasonality refers to a recurring cyclical trend that occurs at consistent intervals. The study of seasonality and temporal trends in crime date back to the mid-19th century and remains a topic of research today [4]. Assessing the seasonal variations in crime patterns has theoretical and policy implications. Different seasons have been reported to be associated with different kinds of crime [5]. The study of the seasonality of crime can help identify the periodicity of crimes and the underlying factors

behind various crimes. Moreover, it can also be utilized in forecasting the volume of crimes to reach an informed decision for taking offence preventive measures and activities required in each season [6]. Studies have shown seasonal fluctuations in sexual violence-related cases [7, 8]. It has also been reported that seasonal patterns of crime can be different between various regions and even between different parts of one country [6, 9]. The nature and frequency of MLCs can vary from region to region, possibly affected by different cultures and seasonal variations [10, 11]. However, the seasonality of sexual assault-related MLCs has not been widely reported in Pakistan.

In the present study, the seasonal decomposition was performed on sexual assault-related MLCs that were conducted in the medicolegal department of Liaquat University Hospital, Hyderabad from 2020 to 2022. The time series analysis method was adopted for seasonal decomposition.

METHODS

This study based on the seasonal trend of sexual assault-related MLCs in Hyderabad region of Pakistan, the relevant data were collected from the casualty department of Liaquat University Hospital, Hyderabad after getting ethical approval from the Institutional Review Board vide letter # LUMHS/FM/44/20, dated: 17-09-2020. The casualty data included the cases that were identified as rape- and sodomy-related cases from January 2020 to December 2022. After arrival of cases of sexual victimization to casualty department, a thorough initial assessment of cases was performed to determine extent of the injury and identify any life-threatening conditions. This involved assessment of vital signs, airway, breathing, and circulation. Any fractures, soft tissue injuries, and other traumatic injuries were provided with treatment for clinical care. In case of unstable or life-threatening conditions, immediate interventions were initiated in order to stabilize the patient, such as controlling bleeding, ensuring adequate oxygenation and ventilation, and addressing any serious injuries. Following prompt trauma care, the subsequent steps included the plan to prioritize addressing sexually transmitted infections, pregnancy concerns, and psychosocial issues. Empiric treatment for sexually transmitted infections was provided, including, ceftriaxone 500 mg IM injection plus azithromycin 1 g PO (single dose) or metronidazole or tinidazole 2 g PO (single dose). Postcoital emergency contraception was offered to female victims without regard to their menstrual cycle. All subjects were excluded if their nature was judged to be non-sexual violence-related. The data were collected and analyzed using MS EXCEL. Seasonal decomposition was performed using Statgraphics Centurion XIX software. The

data e presented as frequencies and their mean ± SD. The seasonal trend and time series analysis were conducted using seasonal decomposition and were presented as plots.


RESULTS

Most of the cases (N=42) were reported in year 2021. The percentage of total sexual assault-related cases occurring in different months of the year shows that a substantial percentage of such cases were aggregated in mid-year, i.e., June, July, and August (10.86% to 14.13%). The highest numbers of sexual assault-related MLCs were performed in July (Table 1).

Table 1: Month-Wise Frequency of Sexual Assault-Related MLCs for Three Years

Month	Year			Mean ± SD	Total
	2020	2021	2022		
January	1	1	4	2 ± 1.41	6 (6.52%)
February	2	3	1	2 ± 0.81	6 (6.52%)
March	1	1	2	1 ± 0.47	4 (4.34%)
April	1	4	3	2 ± 1.24	8 (8.69%)
May	0	4	2	2 ± 1.63	6 (6.52%)
June	1	6	4	3 ± 2.05	11 (11.95%)
July	3	6	4	4 ± 1.24	13 (14.13%)
August	2	6	2	3 ± 1.88	10 (10.86%)
September	3	1	1	1 ± 0.94	5 (5.43%)
October	1	5	3	3 ± 1.63	9 (9.78%)
November	3	4	3	3 ± 0.47	10 (10.86%)
December	1	1	2	1 ± 0.47	4 (4.34%)
Total	19	42	31	42 ± 31	92

The trend of sexual assault-related MLCs that can be interpreted from data in Table 1 was also reflected by seasonal index plots of rape, sodomy, and total sexual-assault MLCs. As depicted in Figure 1, the sexual assault-related MLCs peaked in the third quarter of the year compared to other months. The seasonal time plot (Fig. 1) also shows that the incidence of rape assault peaks in July, consistent with data in Table 1. The seasonal plot of sodomy indicates a high frequency of sodomy in May; however, this is again followed by a higher peak in July.






Figure 1: Seasonal Index Plots for Rape, Sodomy, and Total Sexual Assault-Related MLCs

Smoothed time series plots of rape, sodomy, and total sexual assault-related MLCs are presented in Figure 2. The smoothed time series plot of sexual assault-related MLCs shows three peaks at the start of the second half of each of the three years. The graph for rape-related MLCs shows a similar trend to total sexual assault-related MLCs. However, the trend of sodomy varies slightly in comparison to rape and total sexual assault-related MLCs. This suggests that rape might dominate the seasonal trend of sexual assault-related MLCs. On the other hand, sodomy might present an additional peak of the seasonal trend with reference to the trend of rape-related assaults.





Figure 2: Smoothed Time Series Plot for Rape, Sodomy, and Total Sexual Assault-Related MLCs

DISCUSSION

Any society that upholds the principles of the rule of law and justice considers sexual violence a criminal act. Many types of crimes are periodic as they exhibit seasonal fluctuations in their volume. Often referred to as crime seasonality, this tendency can be captured in the form of time-series data with a recurring cycle [6]. The present study showed the seasonal fluctuation of sexual violence-related MLCs in the Hyderabad region of Pakistan. The data revealed that July, the hottest month in the region in terms of climatic temperature, presented the highest peaks of sexual assault-related MLCs in the region. This is different than a study from mid-Atlantic region of the United States, which observed the least cases of sexual assault reported to the emergency department in July [7]. An eight-year study in Tunisia reported spring as the most vulnerable season for sexual assault, followed by summer [2]. The findings of the present study are similar to some studies from other regions. Summer was found to be a vulnerable season for sexual assault in Qalyubia, Egypt, with more than 50% of 5544 sexual assaults occurring in the summer season [8]. A study from South Korea regarding the experiences of female sexual assault centres in the Incheon metropolitan city found July and August as relatively vulnerable months for sexual assaults [12]. Another study from a different town in South Korea also reported summer as the relatively vulnerable season for sexual assaults [13]. The highest trend of sexual assaults in summer was also found in three Regions of the Republic of Bulgaria [14]. In an Indian tertiary care hospital, the appraisal of sexual offence cases showed the occurrence of most cases during summer season [15]. It has been reported that the same type of crime can exhibit varying seasonal patterns between different countries and regions [9]. Regional differences can occur even within the same nation [6]. Alam et al., observed a major frequency of sexual assaults in Peshawar, Pakistan, occurring in winter in contrast to the present study's findings [16]. However, the number of sexual assault-related MLCs and assessment of their frequencies with reference to seasons has not been reported widely in Pakistan, suggesting the need for more studies. The peaks in violent crimes have

been observed usually in relation to the summer season [17]. This may be ascribed to a reduction in people's mental capacity during warm weather, leading to making of irrational decisions [6]. One of the earliest discoveries in the field of criminology focused on the connection between warm weather and violent behavior. In 18th century, studies were conducted that identified a significant link between high temperatures and crimes against individuals. Researchers also named even named this phenomenon "the thermic law of delinquency" due to its consistent nature. Over the nearly two centuries since then, this foundational correlation has been extensively examined, questioned, and nuanced through a wide range of research studies [18]. Regarding the influence of weather, it has been stated that incidents of property crimes tended to be higher during winter compared to summer. Conversely, crimes against individuals appeared to be more common in summer, attributed to the warmer temperatures and heightened human emotions, while sexual offenses peaked in spring, a season associated with breeding [5]. Similar to the present study, a plenty of research conducted previously has generally identified peaks in summer for overall offenses like property crimes, violent crimes, and sex crimes [4]. A study in seven major US cities, demonstrated that every 5°C increase in daily mean temperature was associated with a 4.5% rise in sex offenses within the following 0–8 days. These associations were more pronounced during hot and cold seasons compared to moderate seasons and could be amplified by higher relative humidity and precipitation. The links were statistically significant for sodomy, fondling, and rape, particularly in specific locations such as open spaces, educational institutions, and streets rather than residences [19]. A cross-over study across Japan reported a nearly linear increase in the relative risk for self-harm and assault behavior as the temperature rose [20]. Likewise, it has been widely found that violent incidents exhibit a seasonal pattern, with the majority occurring during the summer or hotter seasons rather than in winter [17]. Consequently, interpersonal violence during warm weather is expected to persist and probably escalate in the future with temperatures rise as a result of climatic change. For instance, a study in US estimated estimate that 2.6% of sex offenses can be attributed to temperatures exceeding city-specific median temperatures, equating to an average annual sex offense rate of 2.9 per 100,000 individuals. These results highlighted the potential increase in sexual crimes associated with climate change, offering valuable insights for targeted prevention efforts [19, 20]. Another possible reason is the relationship between temperature climates and the activities of people. People tend to spend their time outside in summer compared to other seasons, making them more vulnerable

by staying in a non-protected environment. Moreover, the school-going children remain on holidays and stay out of school during July and August. Consequently, comparatively increasing people's movement out of the relatively protected environment may result in periodic peaks in sexual assaults and other offences [5]. The strategies to prevent violent and sexual crimes linked to hot weathers are yet unclear. One proposed solution is the implementation of heat-warning systems to notify law enforcement agencies about high temperatures and the potential increase in violence. Help lines and public security awareness programs can also play a significant role. However, the effectiveness of these approaches would require assessment in future research studies. Furthermore, more well-designed studies are required to further explore the seasonal decomposition of sexual assault-related MLCs in Pakistan.

CONCLUSIONS

The seasonal fluctuations in sexual assault-related MLCs were assessed in the present study. The results showed the presence of seasonal changes in sexual assault-related MLCs. Smoothed time series and seasonal indices revealed a trend of peaks in sexual assault-related MLCs in summer, with July being the most vulnerable month.

Authors Contribution

Conceptualization: AR

Methodology: AR, UM, NA, AS, MRS, IB

Formal analysis: UM, NA

Writing-review and editing: AR, ASM, MRS, IB

All authors have read and agreed to the published version of the manuscript.

Conflicts of Interest

The authors declare no conflict of interest.

Source of Funding

The authors received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Treglia M, Pallocci M, Passalacqua P, Sabatelli G, De Luca L, Zanovello C et al. Medico-legal aspects of hospital-acquired infections: 5-years of judgements of the civil court of Rome. *Healthcare*. 2022 Jul; 10(7): 1336. doi: 10.3390/healthcare10071336.
- [2] Brahim O, Turki E, Chebbi E, Fersi O, Fatnassi R. Sexual Assault of Women in the region of Kairouan, Tunisia: an 8-year retrospective study on epidemiological and medicolegal characteristics. *BMC Women's Health*. 2022 Mar; 22(1): 64. doi: 10.1186/s12905-022-01647-8.

- [3] Malik R, Atif I, Rashid F, Abbas M. An analysis of 3105 Medico Legal Cases at Tertiary Care Hospital, Rawalpindi. *Pakistan Journal of Medical Sciences*. 2017 July; 33(4): 926-30. doi: 10.12669/pjms.334.11696.
- [4] Delgado R, Sánchez-Delgado H. The effect of seasonality in predicting the level of crime. A spatial perspective. *PLoS One*. 2023 May 31;18(5):e0285727. doi: 10.1371/journal.pone.0285727.
- [5] Andresen MA and Malleson N. Crime seasonality and its variations across space. *Applied Geography* 2013; 43: 25-35. doi: 10.1016/j.apgeog.2013.06.007.
- [6] Shiode N, Shiode S, Nishi H, Hino K. Seasonal characteristics of crime: an empirical investigation of the temporal fluctuation of the different types of crime in London. *Computational Urban Science*. 2023 May 4;3(1):19. doi: 10.1007/s43762-023-00094-x.
- [7] Tadros A, Sharon MJ, Hoffman SM, Davidov DM. Emergency department visits for sexual assault by emerging adults: is alcohol a factor?. *Western Journal of Emergency Medicine*. 2018 Sep; 19(5): 797. doi: 10.5811/westjem.2018.6.38219.
- [8] El-Din AA, Elkholy SM, Metwally ES, Farag HA. Pattern of female sexual assault in Qalyubia Governorate, Egypt, during the period from 2009 to 2013: a retrospective study. *The American Journal of Forensic Medicine and Pathology*. 2015 Dec; 36(4): 276-84. doi: 10.1097/PAF.0000000000000182.
- [9] Andresen MA and Malleson N. Intra-week spatial-temporal patterns of crime. *Crime Science*. 2015 Dec; 4: 1-1. doi: 10.1186/s40163-015-0024-7.
- [10] Qadir MA. Forensic Survey for Medico-Legal Cases: Real Figures from Liaquat University Hospital Hyderabad, Sindh. *Alcoholism*. 2020; 92: 2-53.
- [11] Miller P, Button K, Taylor N, Coomber K, Baldwin R, Harries T et al. The impact of COVID-19 on trends of violence-related offences in Australia. *Journal of Epidemiology and Global Health*. 2023 Sep; 13(3): 504-16. doi: 10.1007/s44197-023-00131-2.
- [12] Kim GJ. Adolescent sexual assault: experiences at sexual assault clinic. *Korean Journal of Obstetrics and Gynecology*. 2005: 1645-53.
- [13] Kim SK, Park YK, Kang YD. Analysis of sexual assault victims: Based on data from a One-stop Service Center in Chungcheongnam-do. *Korean Journal of Obstetrics & Gynecology*. 2012 Oct; 55(10): 736-44. doi: 10.5468/KJOG.2012.55.10.736.
- [14] Demireva RD, Dimitrova MR, Burulyanova IS, Viktorov DW, Geogiev KY. Comparative Investigations of the Incidence of Sexual Assault in Three Regions of the Republic of Bulgaria during 1997-2006. *Acta Facultatis Medicae Naissensis*. 2013 Apr; 30(2): 85. doi: 10.2478/v10283-012-0041-3.
- [15] Mohanty P and Sethi A. Appraisal of Sexual Offence Cases Attending a Tertiary Care Hospital. *Indian Journals*. 2023 Jul; 22(2): 38-43 doi: 10.5958/0974-083X.2022.00041.3.
- [16] Alam N, Naveed S, Ahmad MS, Aziz I. Prevalance of sexual assault at district Peshawar. *Journal of Medical Sciences*. 2018 Sep; 26(3): 225-8.
- [17] Mahendran R, Xu R, Li S, Guo Y. Interpersonal violence associated with hot weather. *The Lancet Planetary Health*. 2021 Sep; 5(9): e571-2. doi: 10.1016/S2542-5196(21)00210-2.
- [18] Thomas C and Wolff KT. Weird winter weather in the Anthropocene: How volatile temperatures shape violent crime. *Journal of Criminal Justice*. 2023 Jul; 87: 102090. doi: 10.1016/j.jcrimjus.2023.102090.
- [19] Xu R, Xiong X, Abramson MJ, Li S, Guo Y. Association between ambient temperature and sex offense: A case-crossover study in seven large US cities, 2007-2017. *Sustainable Cities and Society*. 2021 Jun; 69: 102828. doi: 10.1016/j.scs.2021.102828.
- [20] Kubo R, Ueda K, Seposo X, Honda A, Takano H. Association between ambient temperature and intentional injuries: a case-crossover analysis using ambulance transport records in Japan. *Science of the total environment*. 2021 Jun; 774: 145511. doi: 10.1016/j.scitotenv.2021.145511.