



Original Article



Evaluation of Use of Intensive Care Unit (ICU) Scoring Systems among Healthcare Professionals

Aneela Jahangir^{1*}, Noor-Us-Saba², Madiha Raza², Furkan Hyder² and Syed Farjad Sultan³

¹Department of General Anesthesia, Hospital Manchester University, National Health Service Foundation Trust, Wythenshawe, United Kingdom

²Department of Anesthesia, Dow University of Health Sciences, Karachi, Pakistan

³Department of Anesthesia, Critical Care and Pain Management, Dow University of Health Sciences, Karachi, Pakistan

ARTICLE INFO

Keywords:

ICU Scoring Systems, Critical Care Prognosis, Severity Assessment, Pakistan Healthcare Evaluation

How to Cite:

Jahangir, A., Saba, N.-U.-., Raza, M., Hyder, F., & Sultan, S. F. (2025). Evaluation of Use of Intensive Care Unit (ICU) Scoring Systems among Healthcare Professionals: Intensive Care Unit Scoring Systems Evaluation. *Pakistan Journal of Health Sciences*, 6(1), 119-123. <https://doi.org/10.54393/pjhs.v6i1.1526>

***Corresponding Author:**

Aneela Jahangir
Department of General Anesthesia, Hospital Manchester University, National Health Service Foundation Trust, Wythenshawe, United Kingdom
aneela-jk@hotmail.com

Received date: 15th April, 2024

Acceptance date: 6th January, 2025

Published date: 31st January, 2025

ABSTRACT

The prognosis of patients in the Intensive Care Unit (ICU) is a matter of worry for critically ill patients, their families, and healthcare personnel. However, predicting the chances of recovery in the ICU can be challenging. Scoring systems serve as standardized instruments utilized in critical care research to determine which patients should be included in a study and to establish the comparability of different patient groups. **Objective:** To determine the frequency of use of Scoring Systems to predict the outcome of critically ill patients in the intensive care units of Pakistan. **Methods:** This cross-sectional study was conducted in all the healthcare institutions of Pakistan that are recognized by CPSP for fellowship in ICU training, from 10th April 2018 till 10th October 2018. Forty-one ICU physicians were included. Pre-designed questionnaires were sent to these physicians. **Results:** Out of 41 physicians, 33 (80.5%) of them used and 8 (19.5%) did not use ICU scoring systems. Only 3 consultants reported the use of Anatomical scoring systems. GCS was the most used Disease Specific Scoring System (97.5%), followed by Child Pugh's Score (80%) and Ranson's Criteria (70%). 50% reported the use of Sequential Organ Failure Assessment (SOFA) and 42% used Multiple Organ Dysfunction Score (MODS). 63% used Acute Physiology and Chronic Health Evaluation (APACHE). 93% did not use any Therapeutic Weighted Score. 77.5% used Richmond Agitation-Sedation Scale (RASS) and 25% used Ramsey's. **Conclusion:** It was found that 80.5% physicians used ICU scoring systems to assess mortality and severity of illness.

INTRODUCTION

Critical illness scoring systems are useful for guiding therapy and evaluating the degree of disease and organ failure [1]. Their usage allows one to forecast the overall performance of an intensive care unit, as well as the patient's clinical results and in-hospital mortality [2]. When making important clinical decisions, these rating systems might be helpful. Consequently, hospital resources may be utilized to their full potential, resulting in a decrease in total cost [3]. One such purpose for them is to sort really sick patients into different groups for clinical studies [4]. Although there isn't a universally accepted way to classify the several ICU scoring systems, they can be grouped according to the system in question, the illness at play, the

organ in question, the degree of physiological disturbance, or the therapeutic procedures that have taken place. In addition to this, there are ratings that evaluate more routine aspects of intensive care unit treatment, such as pain management and sedative depth [5]. Early evaluations of outcome prediction relied on doctors' subjective assessments, which Florence Nightingale initiated in 1863 [2]. The fast evolution of intensive care units necessitated the subsequent establishment of scoring systems. Scores have been utilized extensively to evaluate results in intensive care units on a global scale. Additionally, several scoring methods have been contrasted to ascertain whether one produces superior outcomes. Sixty patients in



critical care were the subjects of a prospective research by Sekulic AD et al [6]. Each patient was given an assessment of APACHE II, SAPS II, and MPM II when they were admitted to the intensive care unit. Multiple time periods following admission were assessed for the SOFA and MPM II: 24 hours, 48 hours, 72 hours, and 7 days later. Consequences were strongly predicted by APACHE II and SAPS II scores. The results of this study led to the introduction of consistent use of APACHE II and SAPS II scores upon ICU admission, as well as MPM II and SOFA scores for the duration of the patient's stay in the ICU. Researchers Hosseini M et al., studied 300 critically sick patients to see how well APACHE II and SOFA predicted outcomes in intensive care units [7]. Those patients who did not make it out of the ICU had far higher APACHE II scores than those who did. Data was collected during the first twenty-four hours of admission for the purpose of comparing LODS and APACHE II in a different study that was conducted on 521 patients who were admitted to the neurological intensive care unit [8]. The death rate that was observed was 10.0%, which is much higher than the mortality rates that were anticipated by LODS and APACHE II, which were 7.2% and 4.8%, respectively. In order to determine whether critically sick patients need sedatives or analgesics, the PAD guidelines advocate using delirium, pain, and sedation score systems in the intensive care unit [9]. Patients on mechanical ventilation are less likely to need ventilator support and spend less time in the Intensive Care Unit (ICU) when pain evaluation is performed, according to research by Haniffa R et al [10]. Various metrics have also been used in Pakistani intensive care units to forecast patient outcomes. Patients in critical illness who had higher initial SOFA scores had a greater death rate, according to research by Akbar A et al [11]. Both Lo ML et al., and Hashmi M et al., came to the same conclusion: a high APACHE-II score was inversely related to the duration of stay and increased death risks [12, 13]. Researchers Naqvi IH et al., found that APACHE II, SOFA, and SAP II scores were greater in the non-survivor group compared to the survivor group [14].]. In another study, three scoring systems including Child-Pugh, APACHE II and III were compared to evaluate their prognostic accuracy for predicting short term mortality in patients diagnosed with liver cirrhosis. Among compared scoring systems, Child-Pugh is found to be most reliable one and APACHE scores were found to be less reliable in hospital settings [15]. Several international surveys have found out how frequently these ICU scoring systems are being used. A UK National Survey conducted by Raffa JD et al., found out that 88% of the ICUs used a sedation scoring system [16]. A European survey published in 2001 also showed that sedation scores are most commonly used in the ICUs of UK and Ireland (the Ramsay scale in 74% of cases) So far, no such survey has been

conducted in Pakistan [17]. Keeping in mind the numerous benefits of these scoring systems, their use should be common in the ICU settings. However, insufficient data exists regarding their use in this country.

Accurate prognosis and severity assessment in intensive care units (ICUs) are essential for improving patient outcomes, optimizing resource allocation, and guiding critical care decisions. Although various ICU scoring systems such as APACHE, SOFA, MODS, and RASS are internationally recognized, there is limited national-level evidence regarding their actual utilization among healthcare professionals in Pakistan. This study addressed the lack of local data by evaluating the frequency, patterns, and types of ICU scoring systems used by Pakistani physicians, aiming to assess current practices and identify gaps in standardized critical care assessment.

METHODS

This cross-sectional study was conducted in all healthcare institutions of Pakistan that are recognized by CPSP for fellowship in ICU training from 10th April 2020 till 10th October 2020 and 41 physicians were enrolled. Before data collection, CPSP letter approved on 23rd of December 2019 with reference number CPSP/REU/ANS-2016-218-1605 was taken. In order to collect the data, Purposive sampling technique was used as study targeted respondents of specific expertise. Sample size was calculated by using following formula

$$n = \frac{\frac{z \cdot p(1-p)}{e^2}}{1 + \frac{z \cdot p(1-p)}{e^2}}$$

Online questionnaire was used to collect data from each hospital which included severity, scoring systems, critically ill, intensive care, risk prediction, APACHE, MODS, RASS, SOFA. Consultants of each ICU in the above-mentioned setting having an ICU working experience of more than 1 year, age greater than 30-70 years, qualification FCPS or any other equivalent foreign degree and either gender were included. Those consultants with an MCPS degree and consultants responding after 6 months were excluded. Pre-designed questionnaires were sent via email to the ICU physicians in Pakistan who are working in institutions recognized for fellowship in ICU training by CPSP. A monthly reminder was sent to the non-responders regularly for 6 months. If they failed to respond within 6 months, they were labeled as "non-responders" in the results. Frequency and percentage were used as categorical variables, whereas Standard Deviation (SD) and mean were utilized as quantitative variables. Data were entered and analyzed using SPSS version 21.0. Descriptive analysis was conducted.

RESULTS

The mean age was 41.25±9.3 years whereas the mean duration of practice was 12.05±9.69 years respectively (table 1).

Table 1: Descriptive Statistics of the Patients(n=41)

Variables	Minimum - Maximum	Mean ± SD
Age (Years)	30 - 63	41.25 ± 9.3
Duration of Practice (Years)	1- 40	12.05 ± 9.69

Out of the total, 33 physicians (80.5%) utilized the ICU scoring system, while 8 (19.5%) did not, with a statistically significant p-value of <0.005. Among the participants, 33 (80.5%) were male, and 8 (19.5%) were female. Additionally, 24 physicians (58.5%) were under 40 years of age, whereas 17 (41.5%) were over 40. Regarding the duration of practice, 25 physicians (61%) had less than 10 years of experience, while 16 (39%) had more than 10 years. Moreover, 38 participants (92.7%) held an FCPS qualification, and 3 (7.3%) had other qualifications (Table 2).

Table 2: Demographic Information of the Patients(n=41)

Variable	N (%)
Yes	33 (80.49%)
No	8 (19.51%)
Male	33 (80.49%)
Female	8 (19.51%)
Age < 40	24 (58.54%)
Age > 40	17 (41.46%)
Practice Duration < 10	25 (60.98%)
Practice Duration > 10	16 (39.02%)
FCPS	38 (92.68%)
Other	3 (7.32%)

GCS was the most used Disease Specific Scoring System (97.5%), followed by Child Pugh's Score (80%) and Ranson's Criteria (70%) (Figure 1).

Disease Specific Scoring Systems




Figure 1: Use of different Scoring Techniques for Disease Diagnosis

50% reported the use of SOFA and 42% used MODS. 63% used APACHE. 93% did not use any Therapeutic Weighted Score. 77.5% used RASS and 25% used Ramsey's. Most commonly used Pain Scale was Visual Analogue Scale (VAS) (60%) followed by Numeric Rating Scale (NRS) (50%), 35% used Confusion Assessment Method for Intensive Care Unit (CAM-ICU) (Table 3).

Table 3: Use of Prediction Scoring Systems

Variables	%
Prediction Scoring Systems	
SOFA	50%
MODS	42%
APACHE	93%
RASS	77.5%
Ramsey's	25%
Therapeutic Weighted Score	
Yes	7%
No	93%
Use of Pain Scale	
VAS	60%
NRS	50%
CAM ICU	
VAS	35%
NRS	65%

DISCUSSION

Common applications of general sickness severity ratings in the intensive care unit include outcome prediction, characterization of disease severity and degree of organ dysfunction, and assessment of resource consumption. The many types of scoring are complementary, not mutually exclusive or competitive. The accuracy of indicators for disease severity and prognosis may be enhanced by their potential synergistic effects. It will be required to update all of these scoring systems when new diagnostic, therapeutic, and prognostic techniques become available and as Intensive Care Unit (ICU) demographics change. While some grading systems are more general and used to all patients in the Intensive Care Unit (ICU), others are more disease-or organ-specific, such as the Glasgow Coma Scale (GCS). Modern critical care, patient demographics, and disease frequency have all undergone dramatic changes, and statistical and computational approaches have also come a long way. In the present study, thirty-three (80.5%) and 08 (19.5%) used and did not use any ICU scoring system. Only 3 consultants reported the use of Anatomical scoring systems. GCS was the most used Disease Specific Scoring System (97.5%), followed by Child Pugh's Score (80%) and Ranson's Criteria (70%). 50% reported the use of SOFA and 42% used MODS. 63% used APACHE. 93% did not use any Therapeutic Weighted Score. 77.5% used RASS and 25% used Ramsey's.

Most commonly used Pain Scale was VAS (60%) followed by NRS (50%). 35% used CAM-ICU. In a study, APACHE III and Child-Pugh scores were assessed for all 282 patients. Upper gastrointestinal hemorrhage (38%), liver failure (21%), hepatorenal syndrome (19%), hepatocellular cancer (4%), and spontaneous bacterial peritonitis (6%), were the prominent reasons of death. Survivors had lower Child-Pugh and APACHE III scores (8.6 ± 2.3 and 58.9 ± 35.1) compared to non-survivors (10.9 ± 2.7 and 87.4 ± 30.3) with a p-value less than 0.001. While Child-Pugh properly identified 67% of cases using discriminant analysis, APACHE III accurately recognized 75% of cases ($p < 0.05$) [18]. Every intensive care unit in the United Kingdom took part in a separate postal survey that was conducted across the entire country. There were 192 answers obtained, which is 63.5% of the total number of 302 units that were contacted. A sedative scoring approach was utilized by responding critical care units at a rate of 88%, with the Ramsey sedative Scale being utilized by 66.4% of those centers. In addition to the fact that the majority of units have recorded sedation standards, an astounding 78% of those units also provide evidence that sedation holding is performed on a daily basis. The length of action, rather than the cost, is the major consideration that should be taken into account while selecting a sedative from among the various available options. In intensive care facilities in the United Kingdom, it is routine practice to adhere to sedation guidelines and to use a sedation score tool [19-21]. Another research surveyed 647 critical care doctors from 16 western European nations. Only 35% of those who took the survey reported ever using propofol, while 63% reported ever using midazolam. Midazolam was chosen over propofol in several countries, including France, Germany, the Netherlands, Norway, and Austria, among others. Opioids such as morphine (33%), fentanyl (33%), and sufentanil (24%), were given most often for the purpose of pain treatment. Only 18% of Austrians used a sedative scale, compared to 72% of Britons and 22% of Irish. The Ramsay scale was widely regarded as the most accurate way to measure sedative levels when it was in use [17, 20]. The study had a relatively small sample size and relied on self-reported questionnaire data, which may introduce response bias and limit generalizability. Its cross-sectional design also prevented evaluation of long-term effectiveness or patient outcomes associated with scoring system use. Future research should involve larger multicenter studies with broader representation and assess barriers to implementation, training needs, and the impact of scoring systems on clinical outcomes to enhance evidence-based ICU practices in Pakistan.

CONCLUSIONS

General sickness severity ratings help ICUs evaluate resource utilization, predict prognosis, and characterize disease and organ failure. Disease-specific grading methods are needed because all grades were established for mixed ICU patient groups, reducing subgroup accuracy. ICU populations fluctuate as new diagnostic, therapeutic, and prognostic methods are developed, thus scoring systems must be updated. Since the scoring systems measure diverse things, we think they should complement one other rather than compete. Despite their purpose, organ dysfunction scores predict outcomes. APACHE and SAPS ratings should predict outcomes. These tools assist doctors and management allocate resources and evaluate performance by accurately evaluating illness severity and prognosis.

Authors' Contribution

Conceptualization: AJ

Methodology: NUS, MR, FH

Formal analysis: FH

Writing and Drafting: NUS, MR, FH, SFS

Review and Editing: NUS, MR, FH, SFS

All authors approved the final manuscript and take responsibility for the integrity of the work

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Vincent JL and Moreno R. Clinical review: scoring systems in the critically ill. *Critical Care*. 2010 Apr; 14: 1-9. doi: 10.1186/cc8866.
- [2] Breslow MJ and Badawi O. Severity scoring in the critically ill: part 1-interpretation and accuracy of outcome prediction scoring systems. *Chest*. 2012 Jan; 141(1): 245-52. doi: 10.1378/chest.11-0330.
- [3] Rapsang AG and Shyam DC. Scoring systems in the intensive care unit: a compendium. *Indian journal of critical care medicine: peer-reviewed, official publication of Indian Society of Critical Care Medicine*. 2014 Apr; 18(4): 220. doi: 10.4103/0972-5229.130573.
- [4] Kramer AA, Zimmerman JE, Knaus WA. Severity of illness and predictive models in society of critical care medicine's first 50 years: a tale of concord and conflict. *Critical Care Medicine*. 2021 May; 49(5): 728-40. doi: 10.1097/CCM.0000000000004924.
- [5] Rimawi RH. 2014 bedside critical care guide. 2014 Jan.
- [6] Sekulic AD, Trpkovic SV, Pavlovic AP, Marinkovic OM,

- Ilic AN. Scoring systems in assessing survival of critically ill ICU patients. *Medical Science Monitor: International Medical Journal of Experimental and Clinical Research*. 2015 Sep; 21:2621. doi:10.12659/MSM.894153.
- [7] Hosseini M and Ramazani J. Evaluation of Acute Physiology and Chronic Health Evaluation II and sequential organ failure assessment scoring systems for prognostication of outcomes among Intensive Care Unit's patients. *Saudi Journal of Anaesthesia*. 2016 Apr; 10(2): 168-73. doi: 10.4103/1658-354X.168817.
- [8] Kim TK and Yoon JR. Comparison of the predictive power of the LODS and APACHE II scoring systems in a neurological intensive care unit. *Journal of International Medical Research*. 2012 Apr; 40(2): 777-86. doi: 10.1177/147323001204000244.
- [9] Devlin JW, Skrobik Y, Gélinas C, Needham DM, Slooter AJ, Pandharipande PP, Watson PL, Weinhouse GL, Nunnally ME, Rochweg B, Balas MC. Clinical practice guidelines for the prevention and management of pain, agitation/sedation, delirium, immobility, and sleep disruption in adult patients in the ICU. *Critical care medicine*. 2018 Sep 1; 46(9):e825-73.
- [10] Haniffa R, Mukaka M, Munasinghe SB, De Silva AP, Jayasinghe KS, Beane A, de Keizer N, Dondorp AM. Simplified prognostic model for critically ill patients in resource limited settings in South Asia. *Critical Care*. 2017 Dec; 21: 1-8. doi: 10.1186/s13054-017-1843-6.
- [11] Akbar A, Shahzadi S, Khurram M, Khar HT. Sofa Score and Outcome: Experience at a Public Sector Hospital Icu: Sofa Score And Outcome at A Hospital ICU. *Pakistan Armed Forces Medical Journal*. 2016 Aug; 66(4): 510-14.
- [12] Lo ML, Huang CC, Hu TH, Chou WC, Chuang LP, Chiang MC et al. Quality assessments of end-of-life care by medical record review for patients dying in intensive care units in Taiwan. *Journal of Pain and Symptom Management*. 2020 Dec; 60(6):1092-9. doi:10.1016/j.jpainsymman.2020.07.002.
- [13] Hashmi M, Asghar A, Rashid S, Hameed F. APACHE II analysis of a surgical intensive care unit population in a tertiary care hospital in Karachi (Pakistan). *Anaesthesia, Pain & Intensive Care*. 2019 Jan: 338-44.
- [14] Naqvi IH, Mahmood K, Ziaullah S, Kashif SM, Sharif A. Better prognostic marker in icu-apache ii, sofa or sap ii!. *Pakistan Journal of Medical Sciences*. 2016 Sep; 32(5): 1146. doi: 10.12669/pjms.325.10080.
- [15] Chatzicostas C, Roussomoustakaki M, Notas G, Vlachonikolis IG, Samonakis D, Romanos J, Vardas E, Kouroumalis EA. A comparison of Child-Pugh, APACHE II and APACHE III scoring systems in predicting hospital mortality of patients with liver cirrhosis. *BMC gastroenterology*. 2003 Dec; 3:1-8.
- [16] Raffa JD, Johnson AE, O'Brien Z, Pollard TJ, Mark RG, Celi LA et al. The global open source severity of illness score (GOSSIS). *Critical Care Medicine*. 2022 Jul; 50(7): 1040-50. doi: 10.1097/CCM.0000000000005518.
- [17] Meade MO and Cook DJ. A critical appraisal and systematic review of illness severity scoring systems in the intensive care unit. *Current Opinion in Critical Care*. 1995 Jun; 1(3): 221-7. doi: 10.1097/00075198-199506000-00012.
- [18] Butt AK, Khan AA, Alam A, Shah SW, Shafqat F, Naqvi AB. Predicting hospital mortality in cirrhotic patients: comparison of Child-Pugh and Acute Physiology, Age and Chronic Health Evaluation (APACHE III) scoring systems. *The American journal of gastroenterology*. 1998 Dec 1; 93(12):2469-75.
- [19] Khan MA, Shahbaz H, Noorali AA, Ehsan AN, Zaki M, Asghar F et al. Disparities in adult critical care resources across Pakistan: findings from a national survey and assessment using a novel scoring system. *Critical Care*. 2022 Jul; 26(1): 209. doi: 10.1186/s13054-022-04046-5.
- [20] Pellathy TP, Pinsky MR, Hravnak M. Intensive care unit scoring systems. *Critical Care Nurse*. 2021 Aug; 41(4): 54-64. doi: 10.4037/ccn2021613.
- [21] Jeong S. Scoring systems for the patients of intensive care unit. *Acute and Critical Care*. 2018 May; 33(2): 102. doi: 10.4266/acc.2018.00185.