



Original Article



Comparison of Hospital Duration and Analysis of Harmonic Scalpel Surgery with Conventional Protocols in Patients Undergoing Thyroid Surgery

Dharmoon Arija¹, Muhammad Ameer Kiani², Muhammad Nadeem³, Allah Nawaz⁴, Ahmed Hassan Khan⁴, Khalid Mahmood⁵ and Rana Hamad Ullah⁶

¹Department of Surgery, Ghulam Muhammad Mahar Medical College, Sukkur, Pakistan

²Department of General Surgery, Sheikh Khalifa Medical City Ajman, Ajman, United Arab Emirates

³Department of General Surgery, Hayat Memorial Hospital, Lahore, Pakistan

⁴Department of General Surgery, Sargodha Medical College, Sargodha, Pakistan

⁵Department of General Surgery, Niazi Medical College Sargodha, Sargodha, Pakistan

⁶Pak Red Crescent Medical & Dental College, Lahore, Pakistan

ARTICLE INFO

Keywords:

Thyroidectomy, Conventional Protocols, Harmonic Scalpel, Ultrasonic

How to Cite:

Arija, D., Kiani, M. A., Nadeem, M., Nawaz, A., Khan, A. H., Mahmood, K., & Hamad Ullah, R. (2024). Comparison of Hospital Duration and Analysis of Harmonic Scalpel Surgery with Conventional Protocols in Patients Undergoing Thyroid Surgery: Comparison of Harmonic and Conventional Thyroidectomy. *Pakistan Journal of Health Sciences*, 5(09). <https://doi.org/10.54393/pjhs.v5i09.1528>

*Corresponding Author:

Dharmoon Arija
Department of Surgery, Ghulam Muhammad Mahar Medical College, Sukkur, Pakistan
dharmooarija71@gmail.com

Received Date: 15th April, 2024

Acceptance Date: 17th September, 2024

Published Date: 30th September, 2024

ABSTRACT

A thyroidectomy is a common surgical procedure, and while traditional methods including electrocautery pose a danger of tissue damage, harmonic scalpels have recently seen a rise in their use for this purpose. **Objective:** To determine whether or not the Harmonic Scalpel is beneficial in dealing with patients who are undergoing thyroidectomy. **Methods:** This quasi-experimental study was performed at Department of Surgery, Ghulam Muhammad Mahar Medical College Sukkur from 1st October 2022 to 31st March 2023 and 60 patients were enrolled and divided into 2 equal groups for conventional and harmonic thyroidectomy. Patients with Basedow's illness or thyroid cancer were included regardless of lymph node dissection. For 5mm blood vessel division, 55kHz ultrasound pulses were employed for coagulation. Cutting and coagulation were done using active blade. A well-structured questionnaire recorded operation length, 24-hour drainage volume, hospital stay, post-operative hypocalcaemia, drain placement, RLN paralysis, and blood loss for each patient. **Results:** There were 78.3% females as compared to the males 21.6%. Less operation time was taken in surgery performed by harmonic scalpel as compared to conventional surgery. Significant difference was observed in intra-operative blood loss in study group. Harmonic scalpel lost 40ml blood whereas traditional surgery lost 124ml. Both research groups had similar post-operative problems. **Conclusions:** Harmonic scalpel thyroidectomy appeared to be the most reliable and feasible method as compared to conventional surgery protocol in relevance to intra-operative blood loss, surgery duration, total volume drainage.

INTRODUCTION

Thyroid surgery is considered as the most common endocrine surgery. In spite of the routine surgical procedure, various complications are associated with it including damage to superior laryngeal nerve/recurrent laryngeal nerve, bleeding, hypothyroidism. Another major complication which observed commonly is intra-operative bleeding which occurred due to presence of several blood vessels and narrow visual field. Bleeding leads to numerous post-operative complications and cause air passageway blockage. For that reason, haemostasis in dry field and dissection of blood vessel is imperative to minimize the

adverse effects of post-operative bleeding [1-4]. Advance surgical methods have been developed to combat the negative outcomes of routine surgical procedures. Harmonic surgery is the latest surgical instrument that can cauterize and cut tissue simultaneously. Ultrasonic waves are mainly used in harmonic scalpel and this energy is converted into mechanical energy for blade activation. Activated blade performs the main function and delivers high-grade frictional force. On the other hand, inactivated upper arm helps in the maintaining the tissue position [5-8]. It helps in minimizing the procedure time, blood loss,



volume of drainage in thoracic surgery, abdominal laparoscopic surgery and parotid surgery [7, 8]. Recent studies have also highlighted the role of harmonic scalpel surgery in reduction of surgical time and postoperative blood loss in thyroidectomy as well. Nowadays, harmonic scalpel surgery is actively used in thyroidectomy [9, 10]. Present study was designed for the comparison of hospital duration and analysis of harmonic scalpel surgery with conventional protocols in patients undergoing thyroid surgery.

METHODS

This quasi-experimental study was performed at Department of Surgery, Ghulam Muhammad Mahar Medical College Sukkur from 1st October 2022 to 31st March 2023 with IRB approval letter No. Gen. Surgery/SMBBMU/98. Patients with Basedow' disease or thyroid malignancy were included irrespective of the presence or absence of lymph nodes dissection. The sample size was calculated by using the WHO sample size calculator with power of test (1- β)=80%, level of significance (α)=5%. The difference in the mean hospital stays in groups ligate and group Harmonic Scalpel was 2.18+0.72 and compared to 3.41+1.12 [11]. The sample size was calculated to be 30+30=60 patients. To adapt it to the circumstances, the non-probability purposive sampling method was tweaked. Those patients who were undergoing lobectomy or total thyroidectomy were included in this study. Those patients having complicated comorbidity or autoimmune disease, patient with Hepatitis B and C positive along with the patient with alcoholic abuse and pregnant ladies were excluded from the study. Patients in group A underwent conventional thyroidectomy performed (inferior middle and superior were tied using silk sutures 3/0, and all other vessels were sutured by 4/0 or electro cauterized) and patients in group B were underwent thyroidectomy with harmonic scalpel ligated all thyroid vessels. A generator, a hand piece, and a blade make up the harmonic scalpel arrangement. A piezoelectric crystal stack, compressed between two metal cylinders, serves as the ultrasonic transducer in the hand piece. Mounting the transducer to the blade allows it to be securely fastened. A microprocessor controls the 110-volt generator, which is a high-frequency switching power source, and pulses AC current to the transducer in the hand piece. The transducer may vibrate at 55.5 kHz, its natural harmonic frequency, thanks to this current. The blade that is most commonly utilized in otolaryngological operations resembles a curved paddle. It has a rough outside radius for captive coagulating and a sharp inner beveled edge for cutting. A well-structured questionnaire was used for recording between 6.1-18%g surgery duration, volume of drainage in 24 hours of surgery, hospital stay duration, drain placement duration, post-operative hypocalcaemia RLN paralysis, and blood loss of individual patient. Post-

operative hypocalcaemia was only noted in those patients who were receiving total thyroidectomy. Statistical analysis was performed by using SPSS version 26.0. Chi-square was applied to compare the following variables: time of operation, intraoperative blood loss in milliliters, volume of fluid drainage in milliliters, and number of days spent in the hospital. Student's t-test and the χ^2 -test were used to analyze the results. Difference was considered as significant by considering p-value <0.05.

RESULTS

There were 78.3% females as compared to the males 21.6%. Forty cases of malignant and 20 benign cases were enrolled and divided equally in harmonic scalpel (HS) and conventional surgery (CS) group. Similarly, 30% of the participants in HS group underwent total thyroidectomy whereas 43% of the patients in CS group underwent total thyroidectomy (Table 1).

Table 1: Demographic Variables in the HS and CS Groups (n=60)

Variables	HS N (%) / Mean \pm SD	CS N (%) / Mean \pm SD
Gender		
Female	24 (80%)	23 (76.7%)
Male	6 (20%)	7 (23.3%)
Age (Years)	51.2 \pm 10.79	56.9 \pm 12.79
BMI (Kg/m ²)	22.9 \pm 3.38	24.1 \pm 3.89
Disease Nature		
Malignancy	20 (66.7%)	20 (66.7%)
Benign	10 (33.3%)	10 (33.3%)
Surgery Extent		
Partial Thyroidectomy	21 (70%)	17 (56.7%)
Total Thyroidectomy	9 (30%)	13 (43.3%)

Less operation time was taken in surgery performed by harmonic scalpel as compared to conventional surgery. Significant difference was observed in intra-operative blood loss in study group. Only 40ml blood was lost during HS whereas 124ml blood was lost during CS group. Overall blood drainage volume was also very less in HS group as compared to CS. No difference in hospital stay duration was observed in both study groups (Table 2).

Table 2: Post-Operative Surgical Complications in Study Groups

Variables	Harmonic Scalpel Mean \pm SD	Conventional Surgery Mean \pm SD	p-Value
Operation Time in Minutes	95.3 \pm 18.4	112.2 \pm 21.0	0.045
Intraoperative Blood Loss (mL)	40.1 \pm 32.7	124.8 \pm 39.6	0.055
Gender			
First 24 Hours Post-Surgery	54.4 \pm 24.5	51 \pm 22.7	0.041
Overall volume	119 \pm 87.8	178 \pm 91.6	0.049
Duration of drain placement (Days)	3 \pm 1.5	4 \pm 1.2	0.82
Hospital duration (Days)	3 \pm 1.7	3 \pm 1.4	0.61

Post-operative surgical complications were also assessed in present study. No significant variation was observed in terms of post-operative complications in either of the study group (Table 3).

Table 3: Postoperative Surgical Complications Between the HS and CS Groups

Variables	HC N (%)	CS N (%)	p- Value
Complications	3 (10%)	4 (13.3%)	0.751
RLN Paralysis			
Temporary	2 (6.6%)	2 (6.6%)	0.652
Permanent	-	-	-
Hypocalcemia			
Temporary	2 (6.6%)	2 (6.6%)	0.652
Permanent	-	-	-
Hemorrhage Requiring Surgery	-	-	-
Surgical-Site Infection	-	-	-

DISCUSSION

Thyroidectomy is a routine surgical procedure which can be done through various ways. Conventional clamp/tie techniques are still being employed, however technological advancement is now shifting the trend towards harmonic scalpel surgery. Various adverse effects are associated with conventional procedure such as drainage volume, long procedure time and post-operative bleeding is appeared to be the major concern [12-14]. In the present study, harmonic scalpel surgery showed better outcome in shortening the procedure duration during thyroidectomy in contrast to conventional protocols. This might be attributable to the reason that harmonic scalpel helps in coagulation, detachment and dissection of tissue in a continuous-operation without need of changing instruments. Another important parameter that should be considered is recurrent laryngeal paralysis. It has been proven that lesser chances of RLN paralysis was associated with harmonic scalpel surgery [15-18]. However, in present study, RLN paralysis was not observed in both study groups. Shorter surgical duration minimizes the risk and chances of infections associated with surgeries. It helps in early hospital discharge and accelerates patient recovery process. Shorter surgical duration also linked with cost effectiveness of the procedure [19, 20]. Present study also highlights that shorter duration was observed in harmonic scalpel group as compared to the other group. Few limitations are also associated with this study. Small sample size and inclusion of both benign and malignant diseases might influence the study outcomes.

CONCLUSIONS

Harmonic scalpel thyroidectomy appeared to be the most reliable and feasible method as compared to conventional surgery protocol in relevance to intra-operative blood loss, surgery duration, total volume drainage. Post-operative complication rate of harmonic scalpel was similar to conventional method.

Authors Contribution

Conceptualization: DA

Methodology: DA

Formal analysis: MAK

Writing, review and editing: MN, AN, AHK, KM, RHU

All authors have read and agreed to the published version of the manuscript.

Conflicts of Interest

The authors declare no conflict of interest.

Source of Funding

The authors received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Luo Y, Li X, Dong J, Sun W. A comparison of surgical outcomes and complications between hemostatic devices for thyroid surgery: a network meta-analysis. *European Archives of Oto-Rhino-Laryngology*. 2017 Mar; 274: 1269-78. doi: 10.1007/s00405-016-4190-3.
- [2] Ahmed T, Tahir M, Khan I, Khan MI, Parveen S. Comparison of Outcome after Thyroid Surgery Using Harmonic versus Conventional Clamp and Tie Method. *Isra Medical Journal*. 2018 Sep; 10(5).
- [3] Xu SQ, Ma Y, Su HW, Cheng JF, Zhou YX. Comparison of the effects of focus harmonic scalpel and conventional haemostasis on parathyroid function in thyroid surgery. *Journal of Ayub Medical College Abbottabad*. 2019 Sep; 31(4): 481-4.
- [4] Soliman Mohamed S, O Alfay M, Sh Zarad M. Harmonic scalpel versus conventional technique in total thyroidectomy (comparative study). *Al-Azhar Medical Journal*. 2021 Jan; 50(1): 511-20. doi: 10.21608/amj.2021.147966.
- [5] Cordon C, Fajardo R, Ramirez J, Herrera MF. A randomized, prospective, parallel group study comparing the Harmonic Scalpel to electrocautery in thyroidectomy. *Surgery*. 2005 Mar; 137(3): 337-41. doi: 10.1016/j.surg.2004.09.011.
- [6] Haq AU, Mahmood T, Kumari S, Afzal A. Harmonic Scalpel Thyroidectomy Versus Conventional Haemostasis Thyroidectomy. *International Surgery Journal*. 2021 Jan; 8(2): 559-62.
- [7] Butskiy O, Chang BA, Luu K, McKenzie RM, Anderson DW. A systematic approach to the recurrent laryngeal nerve dissection at the cricothyroid junction. *Journal of Otolaryngology-Head & Neck Surgery*. 2018 Jan; 47(1): 57. doi: 10.1186/s40463-018-0306-7.
- [8] Hirshoren N, Kaganov K, Weinberger JM, Glaser B, Uziely B, Mizrahi I et al. Thyroidectomy practice after implementation of the 2015 American Thyroid Association guidelines on surgical options for patients with well-differentiated thyroid carcinoma.

- Journal of the American Medical Association Otolaryngology-Head & Neck Surgery. 2018 May; 144(5): 427-32. doi: 10.1001/jamaoto.2018.0042.
- [9] El Sherpiny WY. A comparative study between harmonic scalpel hemostasis and conventional hemostasis in total thyroidectomy. *International Surgery Journal*. 2020 Mar; 7(4): 954-60. doi: 10.18203/2349-2902.isj20201372.
- [10] Buzdar F, Khan MA, Majeed Ullah Buzdar IA, Laghari HU, Ahmad R. Role of Harmonic Scalpel Versus Conventional Haemostasis Among Patients Undergoing Total Thyroidectomy. *International Surgery Journal*. 2020; 31(5).
- [11] Mehmood B, Nwaz T, Khan MA, Javaid MU, Nadar A, Manzoor S. Comparison of Complications of Harmonic Scalpel Versus Conventional Homeostasis in Patients Undergoing Near-Total Thyroidectomy at a Tertiary Care Hospital. *Pakistan Journal of Medical & Health Sciences*. 2022 Oct; 16(08): 726-. doi: 10.53350/pjmhs22168726.
- [12] Sultan HM, Ammar MS, Gaber A, Hagag MG, Abd El Bary AR. Comparative study between Focus harmonic scalpel and conventional hemostasis in open total thyroidectomy. *Menoufia Medical Journal*. 2018 Jan 1; 31(1): 175-80.
- [13] Noori IF and Mohammad MA. Sutureless Focus Harmonic Thyroidectomy Versus Conventional Clamping and Knot-Tying Technique. *Medical Journal of Babylon*. 2018 Oct; 15(4): 300-5. doi: 10.4103/MJBL.MJBL_45_18.
- [14] Singhal A, Baskota DK, Acharya K. Ultrasonic cutting and coagulation device versus conventional diathermy dissection in thyroid surgery: prospective randomized trial. *International Journal of Otorhinolaryngology and Head and Neck Surgery*. 2018 Mar; 4(2): 335-8. doi: 10.18203/issn.2454-5929.ijohns20180696.
- [15] Ali MNM, Aldardeer AA, Ali AMA, Radwan AA. Total thyroidectomy: conventional suture ligation technique versus sutureless techniques. *International Surgery Journal*. 2019 May; 6: 1862-9. doi: 10.18203/2349-2902.isj20192355.
- [16] Cannizzaro MA, Bianco SL, Borzi L, Cavallaro A, Buffone A. The use of FOCUS Harmonic scalpel compared to conventional haemostasis (knot and tie ligation) for thyroid surgery: a prospective randomized study. *Springerplus*. 2014 Dec; 3: 1-5. doi: 10.1186/2193-1801-3-639.
- [17] Basurto-Kuba EO, Robles-Estrada M, Hurtado-López LM, EDM EO, Campos-Castillo C, Zaldivar-Ramirez FR et al. Safety and Cost-Effectiveness in Thyroidectomy Using the HARMONIC Scalpel Compared to Traditional Hemostasis: A Controlled Clinical Assay. *Surgical Technology International*. 2017 Jul; 30: 141-7.
- [18] Mohamed WB and Ahmed AE. Sutureless versus conventional thyroidectomy. *International Surgery Journal*. 2017 Mar; 4(4): 1385-8. doi: 10.18203/2349-2902.isj20171147.
- [19] Anandaravi BN, Aslam MA, Nair PP. Prospective randomised study using focus harmonic scalpel versus conventional hemostasis for vessel ligation in open thyroid surgery. *International Surgery Journal*. 2017 Mar; 4(4): 1431-7. doi: 10.18203/2349-2902.isj20171156.
- [20] Arora RD, Prajwal SD, Rao KN, Singh A, Nagarkar NM, Abishek AP. A comparison of harmonic scalpel and conventional techniques for Thyroidectomy. *Indian Journal of Otolaryngology and Head & Neck Surgery*. 2023 Dec; 75(4): 3410-4. doi: 10.1007/s12070-023-04002-x.