



## Original Article



## Adverse Perinatal Outcomes: Their Association with Maternal Anemia

Amna Aziz<sup>1</sup>, Shahida Parveen<sup>2</sup>, Bushra Khan<sup>3\*</sup>, Hajra Sultana<sup>1</sup>, Asma Akhtar<sup>1</sup> and Kanwal Raza<sup>1</sup><sup>1</sup>Department of Obstetrics and Gynecology, Nishtar Medical University, Multan, Pakistan<sup>2</sup>Department of Obstetrics and Gynecology, Quaid-e-Azam Medical College, Bahawalpur, Pakistan<sup>3</sup>Department of Obstetrics and Gynecology, Bakhtawar Amin Medical and Dental College, Multan, Pakistan

## ARTICLE INFO

**Keywords:**

Anemia, Pregnancy, Perinatal outcomes, Hemoglobin, Iron supplementation

**How to Cite:**Aziz, A., Parveen, S., Khan, B., Sultana, H., Akhtar, A., & Raza, K. (2025). Adverse Perinatal Outcomes: Their Association with Maternal Anemia : Perinatal Outcomes with Maternal Anemia. *Pakistan Journal of Health Sciences*, 6(1), 107-111. <https://doi.org/10.54393/pjhs.v6i1.2109>**\*Corresponding Author:**

Bushra Khan

Department of Obstetrics and Gynecology, Bakhtawar Amin Medical and Dental College, Multan, Pakistan

bushrakhan@yahoo.com

Received Date: 27<sup>th</sup> September, 2024Acceptance Date: 19<sup>th</sup> January, 2025Published Date: 31<sup>st</sup> January, 2025

## ABSTRACT

The incidence of anemia during the third trimester of pregnancy correlates with a heightened risk of negative postnatal outcomes. **Objective:** To access the relationship between hemoglobin levels in laboring women and perinatal outcome. **Methods:** A prospective cohort study was done at Department of Obstetrics and Gynecology, Nishtar Hospital, Multan from September 2020 to December 2021 using non-probability purposive sampling technique. A total of 135 anemic pregnant women (Group A) fulfilling inclusion and exclusion criteria was recruited for the study from labour room plus 135 non-anemic women were also recruited as control group (Group B). The data was recorded and explored. **Results:** 135 anemic laboring pregnant ladies were taken in this study having mean age  $25.63 \pm 4.53$  years (range; 19-37 years) and 85.2% had ages less than 30 years. Cesarean section deliveries were noted to be in 50.4% in group A (anemic women) versus 48.1% in group B (non-anemic women). Preterm birth was noted to be 29.6% in group A versus 6.7% in group B. Small for gestational age was 53.3% in group A versus 12.6% in group B. Low birth weight was 54.8% in group A versus 16.3% in group B. Still birth was 7.4% only in anemic women while it was not observed in control group. **Conclusions:** The results of this study supported strong relationship of low haemoglobin in mothers with untoward perinatal outcome; growth retarded babies, preterm, Low birth weight and cesarean section rate. Strong media awareness campaigns should be launched for awareness for iron supplementation among targeted population.

## INTRODUCTION

The World Health Organization (WHO) has estimated that around 2 billion world population or approximately one third of the people are affected with anemia and recent estimated have shown great variation in its burden among pregnant women. Maternal anemia is more prevalent in developing countries particularly Africa and Southeast Asia, where it can be encountered in as many as 35-75% of total pregnancies while it is less than 20% in developed nations [1, 2]. The sustainable development goals take very clear steps towards nutritional status that includes the World Health Assembly targets for scaling down anemia in reproductive age groups (15-49 years) to half by 2025 [3, 4]. The threshold up to which anemia in pregnant women can affect perinatal outcomes still inconclusive due to certain issues such as; insufficient biological factors, social and

economic conditions, different environmental conditions and changes in lifestyle modifications as well as prior to or during pregnancy may be implicated in high burdens of low birth weight [5]. Various dietary habits, such as taking low nutritional diets and lower weight gains in pregnancy, are contributing factors towards low intake of the nutrients required which are measured as necessary for the growth of the fetus for example vitamin B and iron. Ionic iron is the essential mineral which may promote the synthesis of new hemoglobin molecules as well as it is the major basis for energy and required in transporting the oxygen to various organs of the body [6, 7]. Lower concentrations of hemoglobin (Hb%) may promote various changes in angiogenesis of placenta which may restrict supply of oxygen thus limiting fetal growth. This may consequently



lead to certain adverse events such as intrauterine growth restriction and may lead to low birth weight [8, 9]. Recent studies have demonstrated that anemia in pregnancy i.e. Hb levels less than 11g/dl is noticeably associated with birth weight less than gestational age in babies in comparison with those ladies having normal hemoglobin levels during pregnancy [10, 11].

Despite the well-documented global burden of maternal anemia, there remains a significant research gap in low-to-middle income countries like Pakistan regarding the precise hemoglobin thresholds at which perinatal outcomes deteriorate, particularly across different trimesters, parity groups, and socioeconomic strata in community rather than hospital settings. The problem is further compounded by low antenatal care attendance, widespread illiteracy, and poor nutritional awareness, as evidenced by only 19.3% of women maintaining regular antenatal visits in this study. Therefore, this study aimed to assess the relationship between hemoglobin levels in laboring women and perinatal outcomes, generating evidence to inform targeted public health interventions and iron supplementation policies for pregnant women in resource-limited regions of Pakistan.

## METHODS

It was a prospective cohort study, carried out at Department of Obstetrics and Gynecology, Nishtar Hospital, Multan from September 2020 to December 2021 using non-probability purposive sampling technique. The study was approved by the Institutional Ethical Review Board of Nishtar Medical University; Multan vide Reference No. 14417-27/NMU. Iron deficiency anemia in third trimester was defined as Hemoglobin <11g/dl and serum ferritin < 30 microgram/liter according to WHO. Inclusion criteria were laboring mothers with maternal age 18-45 years, primigravida or multi gravida (up to 3) and singleton pregnancy accessed by ultrasound. All women selected were otherwise having uncomplicated pregnancies with body mass index of 18.5-24.9kg/m<sup>2</sup>. Exclusion criteria were laboring mothers with multiple pregnancies, medical disorders like diabetes, hypothyroidism, hyperthyroidism, Hypertension, Ischemic heart disease, valvular heart disease, previous history of preterm births and malignancy were precluded from the study. Babies born with neonatal sepsis, birth asphyxia and meconium aspiration were excluded. Sample size was 270 pregnant women calculated by following formula.  $n = z^2 pq/d^2$ , where,  $p = 75\%$  (hypothesized frequency of need of NICU admission in newborn of anemic women),  $q = 100 - p$ ,  $d = 5.2\%$  margin of error,  $z = 1.96$  (95% confidence) [12]. Sample was divided into two equal groups of anemic (Group A) and non-anemic women (Group B). Women were recruited for the study from labour room after informed consent and was assured of

their confidentiality. Detail history was taken from laboring mothers regarding number of children before current pregnancy, inter birth interval, menstrual irregularities, pregnancy termination. Small for Gestational Age (SGA), Low Birth Weight (LBW), and stillbirths, were measured by comparing the frequency of these outcomes in the anemic group (Group A) with the non-anemic group (Group B). The reference values for SGA and LBW were based on the World Health Organization's criteria for gestational age-specific growth percentiles and weight thresholds, respectively. Stillbirths were recorded based on the absence of signs of life at birth, as per the hospital's standard clinical definitions. The data were recorded and explored by using SPSS version 27.0. Descriptive statistics was used to summarize data. Maternal outcomes such as mode of delivery and preterm labour was analyzed in both groups. Fetal outcomes like small for gestational age, low birth weight, still birth, and NICU admission were studied in the anemic and non-anemic group. The Hb% levels of women during labour and birth weights were extracted from the labour room record. Mean or median with respective measures of dispersion was calculated for quantitative variables like age, gestational age of patient. Frequencies and percentages were calculated for categorical variables like maternal anemia and birth weight of baby. Effect modifiers like age, gestational age and parity was controlled by stratification. Post stratification chi-square test was applied to see their effect on outcome. All tests were two sided and judged statistically significant at  $p < 0.05$ .

## RESULTS

A total of 135 anemic laboring pregnant ladies were taken in this study having mean age  $25.63 \pm 4.53$  years (range; 19-37 years) and 115 (85.2%) had ages less than 30 years. Of these 135 laboring anemic women, 53 (39.3%) belonged to rural areas and 82 (60.7%) belonged to urban areas. Majority of these women were poor 90 (66.7%) and 45 (33.3%) were middle income. Similarly, 97 (71.9%) were illiterate while 117 (86.7%) were house-wives and 79 (58.5%) were living in joint family system. Only 26 (19.3%) were having regular antenatal monthly visits to the hospital. Overall Hb level was  $10.04 \pm 0.46$  g/dl in anemic women and  $11.57 \pm 0.54$  g/dl in non-anemic women. Maternal outcome like cesarean section deliveries were noted to be in 50.4% in group A versus 48.1% in group B ( $P = 0.715$ ). Preterm labour resulting in preterm delivery was noted to be 29.6% in group A versus 6.7% in group B ( $P = 0.001$ ). Small for gestational age was 53.3% in group A versus 12.6% in group B ( $P = 0.001$ ). Low birth weight was 54.8% in group A versus 16.3% in group B ( $P = 0.001$ ). Still birth was 7.4% only in anemic women while it was not observed in control group. NICU admission was 67% among anemic women. Early neonatal deaths were never reported in any group.

**Table 1:** Association of Maternal Hemoglobin Levels with Maternal Outcomes

Maternal Outcomes	Hemoglobin Levels (gm/dL)		p-Value
	Group A Mean $\pm$ SD	Group B Mean $\pm$ SD	
<b>Mode of Delivery (Cesarean Deliveries)</b>			
Yes	9.64 $\pm$ 0.36	11.23 $\pm$ 0.77	0.001
No	10.97 $\pm$ 0.88	11.84 $\pm$ 0.23	
<b>Preterm Labour</b>			
Yes	9.32 $\pm$ 0.42	11.51 $\pm$ 0.32	0.001
No	10.98 $\pm$ 0.97	11.91 $\pm$ 0.63	

The association between maternal hemoglobin levels and neonatal outcomes is presented in Table 2. For neonates classified as Small for Gestational Age (SGA), the mean hemoglobin level was significantly lower in anemic mothers (Group A: 9.64  $\pm$  0.41) compared to non-anemic mothers (Group B: 11.31  $\pm$  0.42,  $p = 0.001$ ). Among non-SGA neonates, the hemoglobin levels were higher in both groups, with Group A at 10.77  $\pm$  0.88 and Group B at 11.88  $\pm$  0.71. Similarly, Low Birth Weight (LBW) neonates were more prevalent in Group A, with a mean hemoglobin level of 9.74  $\pm$  0.46 compared to 11.51  $\pm$  0.49 in Group B ( $p = 0.001$ ). For non-LBW neonates, hemoglobin levels in Group A and Group B were 10.90  $\pm$  0.70 and 11.71  $\pm$  0.67, respectively. Stillbirths were reported exclusively in Group A, with a mean hemoglobin level of 9.52  $\pm$  0.37 ( $p = 0.001$ ), while no cases were observed in Group B. These findings underscore the significant impact of maternal anemia on adverse neonatal outcomes, including SGA, LBW, and stillbirths.

**Table 2:** Association of Maternal Hemoglobin Levels with Neonatal Outcomes

Neonatal Outcome	Group A Mean $\pm$ SD	Group B Mean $\pm$ SD	p-Value
<b>Small For Gestational Age</b>			
Yes	9.64 $\pm$ 0.41	11.31 $\pm$ 0.42	0.001
No	10.77 $\pm$ 0.88	11.88 $\pm$ 0.71	
<b>Low Birth Weight</b>			
Yes	9.74 $\pm$ 0.46	11.51 $\pm$ 0.49	0.001
No	10.90 $\pm$ 0.70	11.71 $\pm$ 0.67	
<b>Still Births</b>			
Yes	9.52 $\pm$ 0.37	NA	0.001
No	10.85 $\pm$ 0.89	NA	

## DISCUSSION

Maternal anemia still remains global health issue with most of the burden falls among developing countries such as Africa, Southeast Asia and in this country, Pakistan [13]. Recent estimates have demonstrated that these countries harbor maternal anemia as much as double high in comparison with developed countries. Data have shown maternal anemia may affect 35-75% of pregnancies developing countries while this figure drops to be 19% for developed countries. It was well narrated that maternal anemia was a powerful predictor for poor perinatal outcomes (both maternal and neonatal) such as low birth

weight, preterm deliveries, NICU admissions, small for gestational age and still births [14]. Furthermore, recent reports from developing nations have also implicated maternal mortality was two times high among severely anemic pregnant women [15]. A total of 135 anemic laboring pregnant ladies were taken in this study having mean age 25.63  $\pm$  4.53 years (range; 19-37 years) and 115 (85.2%) had ages less than 30 years. Debella A et al., from Ethiopia has also reported 26.6  $\pm$  6.15 years mean age of the anemic pregnant women, similar to this case series [16]. Mahmood T et al., from Bahawalpur has also reported 29.4  $\pm$  12.5 years mean age of the anemic pregnant ladies, similar to these results [17]. Bone JN et al., from India has reported similar findings among anemic pregnant women [18]. Kabir MA et al., from Bangladesh has also described 25.71 years mean age of the anemic pregnant women, similar to these results [19]. Baig JA et al., has also reported 28.85  $\pm$  5.12 years mean age of the anemic pregnant women, similar to this study results [20]. Of these 135 laboring anemic women, 53 (39.3%) belonged to rural areas and 82 (60.7%) belonged to urban areas. Debella A et al., from Ethiopia has also reported 38.8% anemic pregnant women were from rural areas, like these findings [16]. Kabir MA et al., from Bangladesh has also reported 32% of the anemic pregnant women from rural areas, same as these results [19]. Majority of these women were poor 90 (66.7%) and 45 (33.3%) were middle income. Debella A et al., from Ethiopia has also reported findings like this study [16]. Kabir MA et al., from Bangladesh has also reported majority of the anemic pregnant women were poor, just like these results [19]. Baig JA et al., has also reported same end result [20]. Similarly, 97 (71.9%) were illiterate while 117 (86.7%) were house-wives and 79 (58.5%) were living in joint family system. Only 26 (19.3%) were having regular antenatal monthly visits to the hospital. Debella A et al., from Ethiopia also reported 49.2% of the anemic pregnant women were housewives and 40% were illiterate, similar to these results [16]. Kabir MA et al., from Bangladesh has also reported 89% of the anemic pregnant women were housewives, similar to these results [19]. Baig JA et al., has also reported 64% of anemic pregnant women were living in joint family system, similar to the study results [20]. Cesarean section deliveries were noted to be in 50.4% in group A versus 48.1% in group B ( $P = 0.715$ ). Mahmood T et al., from Bahawalpur has also reported 45% cesarean section rate among anemic pregnant ladies, similar to our results [17]. Preterm birth was noted to be 29.6% in group A versus 6.7% in group B ( $P = 0.001$ ). Debella A et al., from Ethiopia has also reported 22.2% preterm births among anemic pregnant women, similar to our results [16]. Mahmood T et al., from Bahawalpur has also reported 39% preterm birth among anemic pregnant ladies versus 15% among control group ( $P = 0.001$ ), similar to our results [17]. Kabir MA et al., from

Bangladesh has also reported 40.5% preterm births among anemic pregnant women, similar to our results [19]. Baig JA et al., has also reported 36% preterm births among anemic pregnant women compared with 18% in non-anemic group (P=0.001), similar to these study results [20]. Low birth weight was 54.8% in group A versus 16.3% in group B (P = 0.001). Debella A et al., from Ethiopia has also reported 18.8% low birth weight among anemic pregnant women, lower than our results [16]. Mahmood T et al., from Bahawalpur has also reported 59% low birth weight among anemic pregnant ladies versus 23% in control group (P=0.001), similar to our results [17]. Kabir MA et al., from Bangladesh has reported 18.3% low birth weight among anemic pregnant women [19]. Baig JA et al., has also reported 63% low birth weight among anemic pregnant women versus 18% in non-anemic group, similar to our study results [20]. Small for gestational age was 53.3% in group A versus 12.6% in group B (P = 0.001). Mahmood T et al., from Bahawalpur has also reported 73% small for gestational age among anemic pregnant ladies versus 23% among control group, similar to our results [17]. Still birth was 7.4% only in anemic women while it was not observed in control group. Debella A et al., from Ethiopia has also reported 4.9% still births among anemic pregnant women, just like our results [16]. Mahmood T et al., from Bahawalpur has also observed 8% still births among anemic pregnant ladies versus 3% among control group, same to our results [17]. Another study by Kabir MA et al., in Bangladesh has also stated 8% still births rate among anemic pregnant women, close to our outcome [19]. Baig JA et al., has also mentioned same findings [20].

Although this prospective cohort study provides valuable insights into the association between maternal anemia and adverse perinatal outcomes in Southern Punjab, it carries several limitations, including a relatively small sample size of 270 women, restriction to a single tertiary care hospital limiting generalizability, use of purposive non-probability sampling introducing selection bias, and exclusion of women with comorbidities which may underrepresent real-world complexity. Future studies should employ larger, multi-center, community-based designs across diverse socioeconomic settings in Pakistan, incorporating longitudinal follow-up of neonatal outcomes and evaluating the effectiveness of iron supplementation programs and nutritional interventions on reducing anemia-related perinatal morbidity.

## CONCLUSIONS

These results supported that strong relationship of maternal anemia with deleterious perinatal sequel. Low birth weight, growth retarded babies, cesarean section rate and preterm births were the major adverse outcomes observed in our study. Mean hemoglobin level was significantly lower among anemic pregnant women with

adverse events as compared with those having normal pregnancy outcomes. Strong media awareness campaigns should be launched at national level for awareness for iron supplementation among targeted population to overcome these adverse events.

## Authors' Contribution

Conceptualization: AA<sup>1</sup>

Methodology: BK, HS, AA<sup>2</sup>, KR

Formal analysis: BK, HS

Writing and Drafting: AA<sup>1</sup>, SP

Review and Editing: AA<sup>1</sup>, SP, BK, HS, AA<sup>2</sup>, KR

All authors approved the final manuscript and take responsibility for the integrity of the work

## Conflicts of Interest

The authors declare no conflict of interest.

## Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

## REFERENCES

- [1] Rahman MM, Abe SK, Rahman MS, Kanda M, Narita S, Bilano V et al. Maternal anemia and risk of adverse birth and health outcomes in low-and middle-income countries: systematic review and meta-analysis. *The American Journal of Clinical Nutrition*. 2016 Feb; 103(2): 495-504. doi: 10.3945/ajcn.115.107896.
- [2] Barut A and Mohamud DO. The association of maternal anaemia with adverse maternal and foetal outcomes in Somali women: a prospective study. *BioMed Central Women's Health*. 2023 Apr; 23(1): 193. doi: 10.1186/s12905-023-02382-4.
- [3] Pathirathna ML, Samarasekara BP, Dasanayake TS, Saravanakumar P, Weerasekara I. Adverse perinatal outcomes in COVID-19 infected pregnant women: a systematic review and meta-analysis. *InHealthcare*. 2022 Jan; 10(2): 203. doi: 10.3390/healthcare10020203.
- [4] Getaneh T, Asres A, Hiyaru T, Lake S. Adverse perinatal outcomes and its associated factors among adult and advanced maternal age pregnancy in Northwest Ethiopia. *Scientific Reports*. 2021 Jul; 11(1): 14072. doi: 10.1038/s41598-021-93613-x.
- [5] Kharate MA, Choudhari SG. Effects of Maternal Anemia Affecting Fetal Outcomes: A Narrative Review. *Cureus*. 2024 Jul; 16(7). doi: 10.7759/cureus.64800.
- [6] Detlefs SE, Jochum MD, Salmanian B, McKinney JR, Aagaard KM. The impact of response to iron therapy on maternal and neonatal outcomes among pregnant women with anemia. *American journal of obstetrics*

- & gynecology MFM. 2022 Mar; 4(2): 100569. doi: 10.1016/j.ajogmf.2022.100569.
- [7] Harrison RK, Lauhon SR, Colvin ZA, McIntosh JJ. Maternal anemia and severe maternal morbidity in a US cohort. *American journal of obstetrics & gynecology MFM*. 2021 Sep; 3(5): 100395. doi: 10.1016/j.ajogmf.2021.100395.
- [8] Parks S, Hoffman MK, Goudar SS, Patel A, Saleem S, Ali SA et al. Maternal anaemia and maternal, fetal, and neonatal outcomes in a prospective cohort study in India and Pakistan. *BJOG: An International Journal of Obstetrics & Gynecology*. 2019 May; 126(6): 737-43. doi: 10.1111/1471-0528.15585.
- [9] Rahman MA, Khan MN, Rahman MM. Maternal anaemia and risk of adverse obstetric and neonatal outcomes in South Asian countries: a systematic review and meta-analysis. *Public Health in Practice*. 2020 Nov; 1: 100021. doi: 10.1016/j.puhip.2020.100021.
- [10] Drukker L, Hants Y, Farkash R, Samueloff A, Granovsky SG. 626: Gestational anemia is an independent risk factor for cesarean delivery and adverse perinatal outcome. *American Journal of Obstetrics & Gynecology*. 2014 Jan; 210(1): S306-7. doi: 10.1016/j.ajog.2013.10.659.
- [11] Tian ML, Ma GJ, Du LY, Xiao YG, Zhang YK, Tang ZJ. Prevalence and adverse perinatal outcomes of anaemia in the third trimester of pregnancy in Hebei Province, China. *International Health*. 2024 Jan; 16(1): 91-6. doi: 10.1093/inthealth/ihad028.
- [12] Fowkes FJ, Moore KA, Opi DH, Simpson JA, Langham F, Stanisic DI et al. Iron deficiency during pregnancy is associated with a reduced risk of adverse birth outcomes in a malaria-endemic area in a longitudinal cohort study. *BioMed Central Medicine*. 2018 Dec; 16: 1-0. doi: 10.1186/s12916-018-1146-z.
- [13] Chu FC, Shao SS, Lo LM, Hung TH. Association between maternal anemia at admission for delivery and adverse perinatal outcomes. *Journal of the Chinese Medical Association*. 2020 Apr; 83(4): 402-7. doi: 10.1097/JCMA.0000000000000215.
- [14] Young MF, Oaks BM, Rogers HP, Tandon S, Martorell R, Dewey KG et al. Maternal low and high hemoglobin concentrations and associations with adverse maternal and infant health outcomes: an updated global systematic review and meta-analysis. *BioMed Central Pregnancy and Childbirth*. 2023 Apr; 23(1): 264. doi: 10.1186/s12884-023-05489-6.
- [15] Ali SA, Tikmani SS, Saleem S, Patel AB, Hibberd PL, Goudar SS et al. Hemoglobin concentrations and adverse birth outcomes in South Asian pregnant women: findings from a prospective Maternal and Neonatal Health Registry. *Reproductive Health*. 2020 Nov; 17: 1-3. doi: 10.1186/s12978-020-01006-6.
- [16] Debella A, Eyeberu A, Getachew T, Atnafe G, Geda B, Dheresa M. Perinatal outcomes in anemic pregnant women in public hospitals of eastern Ethiopia. *International Health*. 2023 May; 15(3): 274-80. doi: 10.1093/inthealth/ihac021.
- [17] Mahmood T, Rehman AU, Tserenpil G, Siddiqui F, Ahmed M, Siraj F et al. The association between iron-deficiency anemia and adverse pregnancy outcomes: a retrospective report from Pakistan. *Cureus*. 2019 Oct; 11(10). doi: 10.7759/cureus.5854.
- [18] Bone JN, Bellad M, Goudar S, Mallapur A, Charantimath U, Ramadurg U et al. Anemia and adverse outcomes in pregnancy: subgroup analysis of the CLIP cluster-randomized trial in India. *BioMed Central Pregnancy and Childbirth*. 2022 May; 22(1): 407. doi: 10.21203/rs.3.rs-1232789/v1.
- [19] Kabir MA, Rahman MM, Khan MN. Maternal anemia and risk of adverse maternal health and birth outcomes in Bangladesh: A nationwide population-based survey. *PLoS One*. 2022 Dec; 17(12): e0277654. doi: 10.1371/journal.pone.0277654.
- [20] Baig JA, Jamal MM, Jamal J, Musarrat M. To determine the association of maternal anemia with perinatal outcome in tertiary care hospital. *Pakistan Armed Forces Medical Journal*. 2020 Apr; 70(2): 302-07.