



Original Article



Evaluating the Impact of Site of Oral Cancer on Quality of Life of Patients

Syeda Gulrukh Saba Shah¹, Syeda Lalarukh Saba Shah², Amir Hameed³, Zia-Ur-Rehman Khalil⁴, Aasim Masood⁴ and Farhana Jabeen Shah⁵

¹Department of Anatomy, Kabir Institute of Medical Sciences, Peshawar, Pakistan

²Department of Community Dentistry, Lahore Medical and Dental College, Lahore, Pakistan

³Department of Community Dentistry, Peshawar Dental College, Peshawar, Pakistan

⁴Department of Community Dentistry, Khyber Medical University, Kohat, Pakistan

⁵Department of Community Medicine, Kabir Medical College, Peshawar, Pakistan

ARTICLE INFO

Keywords:

Mouth Neoplasms, Oral cavity, Chemotherapy, Quality of life

How to Cite:

Shah, S. G. S., Shah, S. L. S., Hameed, A., Khalil, Z. U. R., Masood, A., & Shah, F. J. (2025). Evaluating the Impact of Site of Oral Cancer on Quality of Life of Patients: Impact of Oral Cancer on the Quality of Life of Patients. *Pakistan Journal of Health Sciences*, 6(1), 102-106. <https://doi.org/10.54393/pjhs.v6i1.2274>

***Corresponding Author:**

Syeda Gulrukh Saba Shah
Department of Anatomy, Kabir Institute of Medical Sciences, Peshawar, Pakistan
gulrukshah41@gmail.com

Received date: 30th November, 2024

Acceptance date: 21st January, 2025

Published date: 31st January, 2025

ABSTRACT

The Oral Health Impact Profile-14 was created to evaluate how oral health disorders affect the life of the patient, though it is widely utilized in patients with cancer, still, there is limited information regarding its measurement specifically related to the impact of cancer sites in a patient. **Objectives:** To evaluate the site of oral cancer and its impact on patients' oral Health-related quality of life. **Methods:** Diagnosed oral cancer patients were employed from the dental college of Peshawar, and 134 individuals were recruited. Patients of oral cancer, aged 20-59 of both genders were interviewed by using a convenience sampling technique. The Patients were asked about age gender, dental visit, site of cancer, treatment modalities, and various questions related to oral health impact profile. Data analysis was performed using SPSS version 26.0. **Results:** The study revealed that the majority of patients were aged 20-29 and predominantly male (67.9%), and most frequently occurs in the buccal mucosa (35.1%). The results revealed significant challenges among participants, with 30.6% experiencing difficulties in speech and 28.4% with a reduced sense of taste. Physical pain was reported by 38.8%, with 54.5% having eating uncomfortably. **Conclusions:** It was concluded that the study emphasized how patients' quality of life regarding oral health is greatly impacted by the location of cancer.

INTRODUCTION

Oral cancer is the general term for any cancer of the mucosa of the lip, palate, gums, floor of the oral cavity, and tongue [1]. Globally, 9.6 million people died due to cancer, and 18.1 million newly reported worldwide [2]. Pakistan is one of the top nations with the highest incidence of mouth cancer. It is the second most common cause of death, which accounted for 11.27% of deaths in 2020 [3]. Research highlights that cancer has become the leading cause of death in 91 out of 172 countries worldwide [4]. Oral cancer affects the tongue, floor of the mouth, buccal mucosa, gingiva, lips, retromolar trigone, and hard palate. Oropharyngeal cancer involves the base of the tongue, soft palate, tonsils, and pharyngeal wall. [5], Human

papillomavirus (HPV), socioeconomic factors, tobacco use, and genetics significantly increase the risk of cancer [1]. Furthermore, risky behaviours include the use of betel nuts, tobacco, alcohol, and exposure to viral infections [6]. There are several treatment options for oral cavity cancers including pharmacologic cancer therapy, radiation therapy (RT), and surgery, either used separately or in combination. Early-stage cancers have high cure rates with surgery or radiation therapy, although radiation therapy better maintains function [7]. In the past decade, greater attention has been given to patients' quality of life (QoL) following cancer treatment. Assessing a patient's QoL provides valuable insights into the wide range of health



challenges faced by the patient [8]. The oral health impact profile (OHIP), initially 49 items, was shortened to Oral Health Impact Profile-14 (OHIP-14), this updated version is a reliable tool for assessing how patients' lives are affected by their oral health since it preserves accuracy, sensitivity, and cross-cultural reliability [9]. Head and neck cancer (HNC) affects various regions of the head and neck, leading to significant physical changes such as alterations in appearance and pain, as well as functional impairments, including difficulties with chewing, swallowing, and speaking. Beyond these physical and functional challenges, HNC also has profound psychosocial consequences, such as anxiety, depression, and reduced social functioning. Treatments like surgery, chemotherapy, and radiotherapy often result in additional adverse effects and long-term complications [10]. Despite the substantial burden of oral cancer in the population, there is limited data on the prevalence of oral cancers at specific sites and their association with patients' overall well-being.

Oral cancer is a highly prevalent malignancy in South Asia and is associated with significant functional, physical, and psychosocial impairment that directly reduces patients' oral health-related quality of life (OHRQoL). Although previous studies have broadly assessed quality of life in head and neck cancer patients, limited evidence exists on how the *specific site of oral cancer* influences different dimensions of OHRQoL. Most available research treats oral cancer as a single entity, overlooking site-specific variations such as buccal mucosa, tongue, or floor of mouth involvement. Therefore, this study aimed to evaluate the impact of different oral cancer sites on patients' OHRQoL using the Oral Health Impact Profile-14 (OHIP-14), and to determine whether cancer location is significantly associated with quality-of-life outcomes.

METHODS

The cross-sectional study was performed from August, 2023 to March, 2024, at the Khyber College of Dentistry in the Oral Maxillofacial Department. The Gandhara University's Ethical Review Board granted ethical permission (No. GU/Ethical Committee/2023/198) for the study. The study population comprised 134 diagnosed oral cancer patients. The inclusion criteria for this study required that participants be diagnosed and treated for oral cancer. Participants needed to be aged 20-59, of any gender and must have provided written consent at the time of data collection. Those who did not fit the requirements for inclusion were eliminated from the research. Utilizing the Open Epi calculator (version 3.01), the sample size was estimated to be 110 individuals, based on a 95% confidence level, 5% precision, and 7.7% prevalence rate of oral cancer [11]. However, it was increased to 134 to account for the possible absence of answers and dropouts. Convenience

sampling was used to recruit patients. The goals, methods, and confidentiality of the study were fully disclosed to the participants. They were encouraged to ask questions for clarification. Data were collected through patient interviews, where questions related to age, gender, dental visits, types of treatment, and sites of oral cancers were asked. The questionnaire assesses seven aspects of quality of life functional limitation, physical pain, psychological discomfort, physical disability, psychological disability, social disability, and handicap using a 5-point Likert scale (0=never, 4=always). The total score, calculated using the additive method, indicates that higher scores reflect a poorer oral health-related quality of life, which was administered in person by the researcher to ensure accuracy. The scoring system classified impact into three categories based on the score. A score between 0 and 9 indicated a poor impact, suggesting a minimal or negligible effect on health. Scores ranging from 10 to 18 represented as average impact and a score greater than 18 denoted as strong impact, highlighting a pronounced effect on health. Interviews were conducted privately to guarantee participant comfort and confidentiality. A pre-validated questionnaire was used and Cronbach's value was 0.98 and corrected item-total correlations ranged from 0.77-0.93 [10]. For data analysis, SPSS version 26.0 was used. Descriptive statistics summarized the data, and the Kolmogorov-Smirnov test assessed normality. The Kruskal-Wallis test was applied to examine the relationship between OHIP-14 scores and oral cancer site, with a p-value of 0.05 or less indicating statistical significance.

RESULTS

Most respondents were aged 20-29 years and were predominantly male (67.9%). The buccal mucosa comprised 35.1% of cases, followed by the floor of the mouth at 21.6% and the hard palate at 17.9%. The majority of patients received surgery, radiation therapy, and chemotherapy, accounting for 26.9% of the treatments (Table 1).

Table 1: Detailed Overview of Patient Demographics

Variables	Categories	Frequency (%)
Age	20-29 Years	49 (36.6%)
	30-39 Years	19 (14.2%)
	40-49 Years	42 (31.3%)
	50-59 Years	24 (17.9%)
Gender	Male	91 (67.9%)
	Female	43 (32.1%)
Site of Oral Cancer	Tongue	21 (15.7%)
	Hard Palate	24 (17.9%)
	Buccal Mucosa	47 (35.1%)
	Floor of Mouth	29 (21.6%)
	Lip	13 (9.7%)

Treatment of the Patient	Chemotherapy	27 (20.1%)
	Surgery	24 (17.9%)
	Surgery+ RT	23 (17.2%)
	Surgery+ RT+ Chemo	36 (26.9%)
	RT + Chemo	24 (17.9%)
Dental Visit	Within 6 Months	11 (8.2%)

	Once in A Year	44 (32.8%)
	Never	79 (59.0%)

The results indicated significant challenges across multiple domains like the participants faced functional limitations, with 30.6% having trouble pronouncing words and 28.4% reporting a worsened sense of taste (Table 2).

Table 2: Frequency of Responses to the OHIP-14 Scale

Characteristics	0 = never	1 = Rarely	2 = Occasionally	3 = Often	4=Very Often
Functional Limitations					
Trouble Pronouncing Any Words	36 (26.9%)	12 (9.0%)	41 (30.6%)	22 (16.4%)	23 (17.2%)
Felt A Sense of Taste Worsened	38 (28.4%)	31 (23.1%)	25 (18.7%)	22 (16.4%)	18 (13.4%)
Physical Pain					
Had A Painful Aching	17 (12.7%)	19 (14.2%)	33 (24.6%)	13 (9.7%)	52 (38.8%)
Uncomfortable to Eat	7 (5.2%)	9 (6.7%)	32 (23.9%)	13 (9.7%)	73 (54.5%)
Psychological Discomfort					
Have Been Self-Conscious	22 (16.4%)	17 (12.7%)	17 (12.7%)	39 (29.1%)	39 (29.1%)
Felt Tense	22 (16.4%)	15 (11.2%)	7 (5.2%)	46 (34.3%)	44 (32.8%)
Physical Disability					
Unsatisfactory to Diet	5 (3.7%)	13 (9.7%)	12 (9.0%)	29 (21.6%)	75 (56.0%)
Had to Interrupt Meals	46 (34.3%)	30 (22.4%)	28 (20.9%)	16 (11.9%)	14 (10.4%)
Psychological Disability					
Difficult to Relax	81 (60.4%)	15 (11.2%)	10 (7.5%)	17 (12.7%)	11 (8.2%)
Embarrassed	86 (64.2%)	28 (20.9%)	9 (6.7%)	3 (2.2%)	8 (6.0%)
Social Disability					
Irritable with Other People	65 (48.5%)	24 (17.9%)	11 (8.2%)	18 (13.4%)	16 (11.9%)
Difficulty Doing Usual Jobs	64 (47.8%)	23 (17.2%)	13 (9.7%)	22 (16.4%)	12 (9.0%)
Handicap					
Life Was Less Satisfying	96 (71.6%)	19 (14.2%)	8 (6.0%)	4 (3.0%)	7 (5.2%)
Unable to Function	59 (44.0%)	27 (20.1%)	17 (12.7%)	16 (11.9%)	15 (11.2%)

The findings showed a significant association ($p < 0.001$) between the cancer site and OHIP-14 scores. Poor OHIP impact was prevalent in tongue and hard palate cancers, while strong impact was most common in the buccal mucosa (52.9%) and floor of mouth (30.6%) cancers (Table 3).

Table 3: Association Between the Site of Cancer and OHIP-14

OHIP-14	Site of Oral Cancer					p-value
	Tongue	Hard Palate	Buccal Mucosa	Floor of Mouth	LIP	
Poor Impact: $0 \leq \text{score} \leq 9$	7 (50.0%)	6 (42.9%)	0	0	1 (7.1%)	<0.001*
Average Impact: $10 \leq \text{score} \leq 18$	7 (20.0%)	11 (31.4%)	2 (5.7%)	3 (8.6%)	12 (34.3%)	
Strong Impact: score > 18	7 (8.2%)	7 (8.2%)	45 (52.9%)	26 (30.6%)	0	

*Fishers Exact Value: <0.001 Shows Significant Associations

DISCUSSION

A study on oral cancer reported that age or sex variations were not significantly different at any of the sites and the most common tumors were located on the anterior tongue, followed by the buccal mucosa [12]. Based on other research, the tongue was the place most frequently afflicted, then the gingiva [13]. In another study, the head and neck cancer group had a mean age of 49.75 ± 13.44 years and the most common cancer cases were aged 35-64 years. Males predominated in both groups (66% in the cancer group, and 70% in controls). Cancer was most common on the tongue (36%) [14]. The OHIP-14 responses

showed that most patients reported difficulty with daily tasks (21%), tension (21%), irritability (22%), worry about dental issues (32%), discomfort while eating (33%), and unsatisfactory diet (35%). After treatment, these responses improved, except for tension [15]. A study conducted on oral cancer revealed the mean index value was 4.64 before treatment which then declined to 4.25 post-treatment, highlighting the impact of treatment on patients' well-being [16]. A cross-sectional study with 133 oral cancer patients using OHIP-14 found over 95% reporting negative impacts on oral health-related quality of

life. The highest impacts were on eating (83.5%), speaking (77.4%), and emotional status (64.7%), with physical pain, functional limitations, and disability being the most affected dimensions [17]. The findings of the current study showed a significant association ($p < 0.001$) between cancer sites and OHIP-14 scores. Poor OHIP impact was observed in tongue and hard palate cancers, while the strong impact was most frequent in the buccal mucosa (52.9%) and floor of mouth (30.6%) cancers. Studies that were previously conducted reported the mean OHIP-14 scores for various tumor sites. Patients with mandibular tumors had the highest mean score (31.60 ± 15.73), indicating the greatest oral health impact, followed by those with loco-regional metastasis (30.33 ± 10.70) while the lip cancers had the lowest mean score (16.00 ± 14.99), suggesting a lesser impact [18]. According to a study conducted, the OHIP scores for various cancers consistently indicated worse outcomes, reflecting significant disturbances in oral health. These disturbances adversely affected patients' quality of life, highlighting the widespread impact of oral complications across different cancer types [19]. Patients with buccal mucosa cancer exhibited a lower quality of life, highlighting the need for regular follow-ups and supportive therapy. Preoperative psychological reassurance was essential for enhancing postoperative adaptation to psychosocial challenges [20].

This study was limited by its single-center design and use of convenience sampling, which may restrict generalizability to broader populations. The cross-sectional nature also limits causal interpretation between cancer site and quality of life outcomes. Self-reported data through interviews may introduce recall or response bias. Future studies should include multicenter longitudinal designs with larger, more diverse samples and consider additional psychosocial and clinical variables to better understand long-term quality of life outcomes in oral cancer patients.

CONCLUSIONS

It was concluded that the location of oral cancer significantly impacts patients' quality of life. Policymakers should enhance outcomes by improving access to specialized dental care, and psychosocial support, and integrating routine quality-of-life assessments into cancer care protocols.

Authors' Contribution

Conceptualization: SGSS

Methodology: SGSS, SLSS, AH, ZURK, AM, FJS

Formal analysis: SLSS

Writing and Drafting: ZURK, SGSS, SLSS, AH

Review and Editing: ZURK, AM, FJS, SGSS, SLSS, AH

All authors approved the final manuscript and take responsibility for the integrity of the work

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Khan MF, Hayhoe RP, Kabir R. Exploring the Risk Factors for Oral Cancer in Pakistan: A Systematic Literature Review. *Dentistry Journal*. 2024 Jan; 12(2): 25. doi: 10.3390/dj12020025.
- [2] Duong HY, Rocuzzo A, Stähli A, Salvi GE, Lang NP, Sculean A. Oral Health-Related Quality of Life of Patients Rehabilitated with Fixed and Removable Implant-Supported Dental Prostheses. *Periodontology 2000*. 2022 Feb; 88(1): 201-37. doi: 10.1111/prd.12419.
- [3] Spanemberg JC, Cardoso JA, Slob EM, López-López J. Quality of Life-Related to Oral Health and Its Impact in Adults. *Journal of Stomatology, Oral and Maxillofacial Surgery*. 2019 Jun; 120(3): 234-9. doi: 10.1016/j.jormas.2019.02.004.
- [4] Gao M, Yan X, Zhao R, Shan Y, Chen Y, Jian F *et al*. Comparison of Pain Perception, Anxiety, and Impacts On Oral Health-Related Quality of Life Between Patients Receiving Clear Aligners and Fixed Appliances During the Initial Stage of Orthodontic Treatment. *European Journal of Orthodontics*. 2021 Jun; 43(3): 353-9. doi: 10.1093/ejo/cjaa037.
- [5] Algudaibi LY, AlMeaigel S, AlQahtani N, Shaheen NA, Aboalela A. Oral and Oropharyngeal Cancer: Knowledge, Attitude and Practices Among Medical and Dental Practitioners. *Cancer Reports*. 2021 Aug; 4(4): e1349. doi: 10.1002/cnr.21349.
- [6] Lin X, Wu X, Gomaa A, Chen J, Wu L, Xie X *et al*. Analysis of Risk Factors for Multiple Primary Oral Squamous Cell Carcinoma: A Cohort Study. *Clinical Oral Investigations*. 2020 Sep; 24: 3147-55. doi: 10.1007/s00784-019-03189-0.
- [7] Kosgallana S, Jayasekara P, Abeysinghe P, Lalloo R. Oral Health-Related Quality of Life of Oral Cancer Patients Treated with Radiotherapy Alone or with Chemotherapy in A Tertiary Referral Centre in Sri Lanka. *BioMed Central Oral Health*. 2023 Mar; 23(1): 162. doi: 10.1186/s12903-023-02854-x.
- [8] Vermaire JA, Partoredjo AS, de Groot RJ, Brand HS, Speksnijder CM. Mastication in Health-Related Quality of Life in Patients Treated for Oral Cancer: A Systematic Review. *European Journal of Cancer Care*. 2022 Nov; 31(6): e13744. doi: 10.1111/ecc.13744.
- [9] Janto M, Iurcov R, Daina CM, Neculoiu DC, Venter AC, Badau D *et al*. Oral Health among Elderly, Impact On

- Life Quality, Access of Elderly Patients to Oral Health Services and Methods to Improve Oral Health: A Narrative Review. *Journal of Personalized Medicine*. 2022 Feb; 12(3): 372. doi: 10.3390/jpm12030372.
- [10] Tesic M, Cankovic M, Jevtic M, Stevanovic D. Validation of the Oral Health Impact Profile-14 in Patients with Head and Neck Cancer. *Medicina Oral, Patologia Oral, Cirugia Bucal*. 2020 Nov; 25(6): e739. doi: 10.4317/medoral.23765.
- [11] Goh V, Hassan FW, Baharin B, Rosli TI. Impact of Psychological States On Periodontitis Severity and Oral Health-Related Quality of Life. *Journal of Oral Science*. 2022; 64(1): 1-5. doi: 10.2334/josnusd.21-0267.
- [12] Wang W, Adeoye J, Thomson P, Choi SW. Statistical profiling of oral cancer and the prediction of outcome. *Journal of Oral Pathology and Medicine*. 2021 Jan; 50(1): 39-46. doi: 10.1111/jop.13110.
- [13] Brennan PA, Dylgjeri F, Coletta RD, Arakeri G, Goodson AM. Surgical Tumour Margins and Their Significance in Oral Squamous Cell Carcinoma. *Journal of Oral Pathology and Medicine*. 2022 Apr; 51(4): 311-4. doi: 10.1111/jop.13276.
- [14] Maqbool SH, Siddique SO, Mehak AY, Ali QA, Asghar SK, Shahid FA. Evaluation of Oral Health-Related Quality of Life in Patients Diagnosed with Head and Neck Cancer Receiving Radiotherapy. *Pakistan Journal of Medical and Health Sciences*. 2021 Nov; 15(11): 3119- 3122. doi: 10.53350/pjmhs2115113119.
- [15] Memon AB, Rahman AA, Channar KA, Zafar MS, Kumar N. Evaluating the Oral-Health-Related Quality of Life of Oral Submucous Fibrosis Patients Before and After Treatment Using the OHIP-14 tool. *International Journal of Environmental Research and Public Health*. 2022 Feb; 19(3): 1821. doi: 10.3390/ijerph19031821.
- [16] Priyadharshini G, Ramalingam K, Ramani P, Krishnan M. Longitudinal Assessment of the Quality of Life in Oral Squamous Cell Carcinoma Patients. *Cureus*. 2024 May; 16(5). doi: 10.7759/cureus.60596.
- [17] Caminha RD, Caldas RJ, Bueno IC, Scaraficci AC, da Silva Santos PS. A Time Frame Evaluation of the Oral Health-Related Quality of Life Among Patients with Head and Neck Cancer. *Oral Oncology Reports*. 2024 Sep; 11: 100593. doi: 10.1016/j.oor.2024.100593.
- [18] Ghorbani Z, Manifar S, Bohloli G, Aghakouchakzadeh A, Mirzaei A. Oral Health-Related Quality of Life in Patients with Oral Squamous Cell Carcinoma: A Case-Control Study. *Dental Research Journal*. 2023 Mar; 20(1): 36. doi: 10.4103/1735-3327.372653.
- [19] Soldera EB, Ortigara GB, Bonzanini LI, Schulz RE, Danesi CC, Antoniazzi RP et al. Clinical and Sociodemographic Factors Associated with Oral Health-Related Quality of Life in Survivors of Head and Neck Cancer. *Head and Neck*. 2020 May; 42(5): 886-97. doi: 10.1002/hed.26063.
- [20] Ramalingam K, Krishnan M, Ramani P, Muthukrishnan A. Quality of Life Assessment with European Organisation for Research and Treatment of Cancer Questionnaire (Head and Neck Module 43) and Its Clinicopathological Correlation among Patients Treated for Oral Squamous Cell Carcinoma: An Exploratory Study. *Cureus*. 2023 Feb; 15(2). doi: 10.7759/cureus.34650.