



Original Article



Patient Satisfaction Level and its Various Determinants in a Tertiary Care Hospital of Peshawar, Pakistan: A Cross-Sectional Study

Muhammad Wasim^{1,2*}, Khalid Rehman³, Majid Paracha¹, Junaid Sarfaraz⁴ and Sahibzada Mahmood Noor¹

¹Department of Dermatology, Lady Reading Hospital, Peshawar, Pakistan

²Department of Community Medicine, Kabir Medical College, Peshawar, Pakistan

³Institute of Public Health and Social Sciences, Khyber Medical University, Peshawar, Pakistan

⁴Department of Cardiac Surgery, Lady Reading Hospital, Peshawar, Pakistan

ARTICLE INFO

Keywords:

Patient Satisfaction, Healthcare Quality, Inpatient Care, Likert Scale

How to Cite:

Wasim, M., Rehman, K., Paracha, M., Sarfaraz, J., & Noor, S. M. (2025). Patient Satisfaction Level and its Various Determinants in a Tertiary Care Hospital of Peshawar, Pakistan: A Cross-Sectional Study: Patient Satisfaction Level in a Tertiary Care Hospital. *Pakistan Journal of Health Sciences*, 6(2), 255-261. <https://doi.org/10.54393/pjhs.v6i2.2689>

***Corresponding Author:**

Muhammad Wasim
Department of Dermatology, Lady Reading Hospital, Peshawar, Pakistan
Department of Community Medicine, Kabir Medical College, Peshawar, Pakistan
iftikharahmad1437@gmail.com

Received date: 27th December, 2024

Acceptance date: 21st February, 2025

Published date: 28th February, 2025

ABSTRACT

Patient satisfaction is a surrogate metric for quality of services provided by healthcare system.

Objective: To assess patient satisfaction and associated factors in a major tertiary care hospital. **Methods:** This cross-sectional study was conducted at the Lady Reading Hospital (LRH), Peshawar, Pakistan. All patients admitted for indoor treatment for at least 2 days were enrolled through multistage sampling technique. Patients admitted to ICU, CCU, psychiatry or oncology wards and COVID-19 positivity were excluded. Patient satisfaction level was assessed with the Patient Satisfaction Questionnaire (PSQ)-18. Likert scale was utilized to evaluate patients' response, where the scale ranged from 1 (complete dissatisfaction) to 5 (complete satisfaction). Patients with cumulative score of ≥ 50 were classified as satisfied. **Results:** Of the total 384 patients enrolled, 199 (~52%) patients were older than 45 years of age, while the female to male ratio was 1.25:1. The number of satisfied patients was significantly higher than dissatisfied patients (313 vs. 71; $p < 0.01$). Highest number of patients were satisfied with the domain D7 (i.e., accessibility and convenience: 72%), followed by domain D6 (i.e., time spent with the doctor: 69%). A minimum number of patients were satisfied with the domain D2 (i.e., technical quality: 53%). Moreover, higher number of patients with age ≥ 45 years (~86%), female gender (~86%), with bachelor education (~100%) and patient admitted to medical and allied wards (~86%) expressed satisfaction. **Conclusion:** Overall, the study showed a high patient satisfaction at LRH, Peshawar.

INTRODUCTION

Patient satisfaction is a vital metric in evaluating services of healthcare system and eliminating claims of malpractice. Studies focused on evaluating patient satisfaction are increasing worldwide [1]. Healthcare providers are always interested in providing cost-effective facilities that can deliver superior and efficient medical services [2]. While a patient is in the hospital, the workflow coordinates both the safety of hospital staff and patients, and the quality of care they receive [3, 4]. To move forward, hospital administrators and healthcare professionals should consider key factors including the quality of patient

care and specialty practices that create competency and capacity. Achieving such goals requires the inclusion of several categories [5]. For example, by analyzing patient data, hospital management can find deeper correlations and data patterns that can lead to improvements in medical procedures and practices [6]. With the pioneering work started in 1950s, patient satisfaction (and improving quality of patient care) is still an ongoing process, and comprise of a combination of technical, interpersonal and organizational aspects [7, 8]. The following six areas have been identified to cover the entire structure of the



healthcare system: Healthcare and medical care should be knowledge-based, safe, patient-centered, effective, equitable and timely [9]. Moreover, many parameters have been related to patient satisfaction, including physician's knowledge, attitudes, and responsiveness, wait times, patient privacy, infrastructure, cleanliness, cost-effectiveness, length of visit, access to medication, counseling and clinical procedures [10]. Patients also differ in their satisfaction with various aspects of the healthcare system, depending on their personal quality of life is influenced by the environmental, social, and informational aspects that affect patient satisfaction with the services availed [11]. Recently, the focus of healthcare system has shifted to the quality of services. In this regard, a simple and direct approach is to ask patients themselves to rate the healthcare system [12]. As a result, the notion of patient satisfaction surveys with questionnaires on the quality of care was established [13]. Patient satisfaction being a key indicator for characterizing the healthcare system and the services it provides to the public, this cross-sectional study was designed to evaluate patient satisfaction and related factors at the Lady Reading Hospital (LRH) Peshawar, Pakistan [14].

Despite increasing emphasis on patient-centered healthcare, limited local evidence exists regarding comprehensive inpatient satisfaction and its determinants in public tertiary care hospitals of Khyber Pakhtunkhwa, particularly using standardized tools like PSQ-18. Previous Pakistani studies have often focused on selective departments or public-private comparisons, leaving gaps in understanding domain-specific satisfaction within large public institutions. Therefore, this study aimed to assess overall patient satisfaction levels and identify demographic, educational, and ward-related factors influencing satisfaction among admitted patients at Lady Reading Hospital, Peshawar.

METHODS

This study was designed as a cross-sectional, single hospital-based study for a duration of 6 months (Jan-June, 2024), was carried at the LRH Peshawar, Khyber Pakhtunkhwa. Patients admitted for indoor treatment at all departments of LRH, Peshawar for at least 2 days were eligible for enrollment in this study. The terms "ward admission" and "indoor treatment" were used interchangeably. All indoor patients were categorized based on their admission into either Surgical and Allied Wards or Medical and Allied Wards. Patients admitted for indoor treatment were included in the study; however, those with specific conditions were excluded to maintain the study's focus on general inpatient satisfaction. Exclusion criteria included patients admitted to the Intensive Care Unit (ICU) or Coronary Care Unit (CCU), as well as those in psychiatry and oncology wards.

Additionally, patients who tested positive for COVID-19 or Congo virus were not enrolled. Pediatric and nursery ward patients were also excluded to ensure the study targeted adult inpatients capable of independently assessing their satisfaction with healthcare services. The OpenEpi Calculator- an open-source statistical tool- was used to estimate the sample size for this study. It was assumed that the data collected in this study will follow the Chi square statistics. Keeping a 50% patients proportion in MTI, statistical significance at 0.05 and a 95% confidence level, the required sample size was estimated, which came out to be 384. After ensuring the eligibility, patients were enrolled through multistage consecutive sampling technique. The Patient Satisfaction Questionnaire (PSQ)-18 was utilized to quantify the level of patient satisfaction [15]. The PSQ-18 consisted of 18 questions, pertaining to the following seven domains: general satisfaction (2 items), interpersonal manner (2 items), time spent with the doctor (2 items), technical excellence (4 items), economical aspects (2 items), accessibility and convenience (4 items). Many items of the PSQ-18 use the term health screening, which is indicative of the particular diagnostic and/or therapeutic services availed at the hospital. The patient response to each of these question was recorded on the Likert scale, ranging from 1 (completely unsatisfied) to 5 (completely satisfied). The Likert scale is a psychometric scale is widely involved in research that employs questionnaires [16]. Thus, the satisfaction level of each patient was reflected by a cumulative score of 18-90. The patient was regarded as dissatisfied if PSQ-18 cumulative score < 50 and satisfied if PSQ-18 cumulative score \geq 50 [17]. The SPSS (version 21) was used for data storing, sorting, manipulation and analysis. Statistical significance was determined using chi-square test. The study was approved by the Ethical Board of Khyber Medical University, Peshawar (Reference No. KMU/IPHSS/Ethics/2023/PS/0171). Patients were enrolled up on agreement for participation, and signing a written consent.

RESULTS

The PSQ-18 consists of two major parts, including patient demographics and patient satisfaction regarding different domains of the healthcare system. Table 1 presented a summary of patient demographics. Of the total 384 patients enrolled, 214 (~ 55.7 %) were female patients. The majority of patients were of old age. Specifically, 199 (~ 51.82%) patients belonged to the age of more than 45 years. There were 58 (~ 15.1%) patients younger than 25 years. The number of patients in 26-25 years and 36-45 years were 63 (16.41 %) and 64 (16.67%), respectively. Education level assessment of the patients revealed that maximum number of patients did not studied at all, followed by the school education (up to grade 10). The number of patients with intermediate, bachelor and graduate level education

were 36 (9.4%), 16 (4.2%) and 02 (0.5%), respectively. The distribution of patients in the medical and allied and surgical and allied was 81.5 and 18.5%, respectively.

Table 1: Demographics of Patients included in this Study

Variables	Category	Frequency (%)
Gender	Male	170 (44%)
	Female	214 (56%)
Age (Years)	< 25	58 (15%)
	26-35	63 (16%)
	36-45	64 (17%)
	>45	199 (52%)
Education	No Education	283 (74%)
	School	47 (12%)
	Intermediate	36 (9.5%)
	Bachelor	16 (04%)
	Master	02 (0.5%)
Ward of Admission	Medical and Allied	313 (81.5%)
	Surgical and Allied	71 (18.5%)

Overall, 82.81% (313/384) of the patients were satisfied with the services and facilities at the hospital, while a relatively small number of patients (71/384: 17.19%) were not satisfied. Table 2 showed stratified data on patient satisfaction in each of the seven domains of PSQ-18. The

Table 3: Satisfaction and Dissatisfaction Frequencies against Each Item of PSQ-18

Questions	Mini Score	Max Score	Median Score	Mean Score	Standard Deviation
Screening received was perfect	1.00	5.00	3	3.21	1.34
Dissatisfied with the screening received	1.00	5.00	3	2.71	1.14
Screening station provided appropriate medical care	1.00	5.00	4	3.77	1.03
I wonder about the correctness of diagnosis	1.00	5.00	3	3.03	1.37
During screening, the staff carefully examined everything	1.00	5.00	4	3.78	1.15
Doubts about the capability of my doctor	1.00	5.00	3	2.89	1.29
Staff acted busy and impersonal	1.00	5.00	3	2.66	0.98
Screening staff was very friendly and courteous	1.00	5.00	4	3.83	1.10
Staff explained the health screening	1.00	5.00	2	2.17	1.16
Doctor sometime ignored me	1.00	5.00	3	3.00	1.31
Screening availed without financial set back	1.00	5.00	3	3.25	1.05
Affordability of health screenings	1.00	5.00	3	2.74	1.13
The screening staff was in too much hurry	1.00	5.00	3	3.08	1.27
Time spent on screening was adequate	1.00	5.00	2	2.27	1.18
Ease in access to the health screenings	1.00	5.00	4	3.59	1.10
Waiting period for the health screening	1.00	5.00	3	2.79	1.14
Getting appointment for health screenings was hard	1.00	5.00	3	2.96	1.05
Health screenings is offered whenever needed	1.00	5.00	3	3.06	1.27

The gender-wise data stratification for patient satisfaction showed significantly higher satisfaction level for female patients compared to male patients (86.5% vs. 78.2%, $p < 0.05$: Figure 1a). Age stratification (Figure 1b) illustrated highest satisfaction rate of 86.4% for older patients of age > 45 years. Focusing on the younger patients, differences in satisfaction rate were seen (<25, 26-35, 36-45 years: 87.9%, 77.8%, 71.9%, $p > 0.05$). However, these differences were not statistically significant ($p > 0.05$). Moreover, it was found that the patients admitted to the medical and allied wards were more satisfied than those admitted to the surgical and allied wards (86.26% vs. 67.61%, $p < 0.01$: Figure 1c). Moreover, the impact of patients' education levels on satisfaction levels was also evaluated (Figures 1d and 2b). Educational attainment was categorized into five levels: No Education, School, Intermediate, Bachelor, and Master. The correlation between education level and satisfaction was

highest number of patients (277: 72%) were satisfied with domain D7 (Accessibility and Convenience), followed by 264 (69%) in the D6 (Time Spent with Doctors). Likewise, 257 (67%) and 248 (65%) patients were satisfied with D5 (Financial Aspects) and D3 (Interpersonal Manner), respectively. The lowest patient satisfaction was observed D2 (202: 53%).

Table 2: Satisfaction and Dissatisfaction Frequencies in Various Domains of PSQ-18

Domain of PSQ-18	Satisfaction Frequency (%)	Dissatisfaction Frequency (%)
D1: General Satisfaction	207 (54%)	177 (46%)
D2: Technical Quality	202 (53%)	182 (47%)
D3: Interpersonal Manner	248 (65%)	136 (35%)
D4: Communication	233 (61%)	151 (39%)
D5: Financial Aspects	257 (67%)	127 (33%)
D6: Time Spent with Doctors	264 (69%)	120 (31%)
D7: Accessibility and Convenience	277 (72%)	107 (28%)

Table 3 presented the minimum, maximum, median and mean Likert score for each of the 18 items of the PSQ-18. Majority of the median and mean scores are equal to or above 3, suggested that majority of the patients were satisfied.

analyzed, with results further stratified by gender to assess variations between male and female patients. Overall, satisfaction among patients with no, school, intermediate, bachelor and master education was 82.69% (234/283), 85.11% (40/47), 75% (27/36), 100% (16/16) and 50% (1/2), respectively (Figure 1d).




Figure 1: Patient satisfaction evaluated using PSQ-18. The number of satisfied patient (x-axis) were stratified on the basis of (a) gender, (b) age, (c) ward admission, and (d) education. White and gray bars represent satisfied and dissatisfied patients, respectively.

Patient satisfaction was further carried out in sub-groups. Results for satisfaction of male and female patients with respect to their age, level of education and ward admission are shown in Figure 2. For the male group, patients older than 45 years of age (87.37%) (Figure 2a), with no or school education (83.3% and 100%) (Figure 2b) and those admitted to medical and allied wards (90.21%) (Figure 2c) showed higher satisfaction. This trend was slightly different in the female patient group. Specifically, the satisfaction rate in female patients older than 45 years of age was 85.58% (Figure 2a), in patients with no or bachelor education was 100% and 82.29% (Figure 2b) while 82.94% in female patients admitted to medical and allied wards (Figure 2c).




Figure 2: Patient satisfaction stratified for male (top row) and female (bottom row) patients. Sub-stratification of male and female patient groups was performed with respect to their age (a), education (b), and ward admission (c). White and gray bars represent satisfied and dissatisfied patients, respectively.

DISCUSSION

The PSQ-18 is a valuable tool that allows to measure patients' satisfaction in a variety of domains [15]. This analyses revealed that the number of satisfied patient was significantly higher than the number of dissatisfied patients (318 vs. 66; $p < 0.01$). Expanding the analysis to each question of PSQ-18 showed that mean satisfaction score in 16/18 question was 60% or above. Nonetheless, the results presented herein may be viewed in the light of their inherent shortcomings. First, a cross-sectional study only provides a snapshot of the prevailing situation. Due to the design of the PSQ-18, the responses may seem subjective. Moreover, although this study enrolled the required number of eligible participants, a multicenter study would provide better understanding of the factors related to patient satisfaction. Such analyses can potentially reveal more heterogeneity and other substructures in the patient experience, leading to a deeper understanding of patient satisfaction and related factors. The quality of care has been divided into the professional quality of the service providers and the functional quality perceived by the patients [18]. Professional quality refers to the level of competence in providing professional medical skills and making accurate diagnoses. On the other hand, patient-perceived quality means not only medical competence but also functional quality, which indicates how well the patient's needs (equipment, facilities, physical environment, communication, etc.) are met [19]. Based on this, it was found that patients with poor medical outcomes were satisfied. In addition, poor quality of care has been reported when patients feel they have not received an accurate diagnosis and treatment [20]. The quality of medical services depends heavily on the criteria that patients use to make their own assessments. As patient needs become more diverse and demanding, subjective assessment from the patient's perspective is considered more important for assessing the quality of medical care [21]. Previous studies investigating patient satisfaction have concluded that the attitude of the medical staff (i.e., doctors, nurses, paramedics) had the highest satisfaction value, while other studies emphasized the importance of trust in the competence and attention of doctors on the quality of care [22]. The existing literature also suggests that the interpersonal communication, soft skills and behaviors of the healthcare provider towards patients are directly related to patient satisfaction [23]. Meanwhile, hospital facilities seemed to be the area that should be given more attention to improve inpatient satisfaction, in line with other findings [24]. Moreover, longer waiting times are negatively correlated with patient satisfaction of the service provider. Studies have also shown that elderly patients, low-income patients, female patients, patients with only primary education and patients in rural areas have high levels of satisfaction [25]. Interestingly, the results of this study are consistent with previous studies on age and

educational level in several developed countries. Beside international studies, several authors have investigated and evaluated patient satisfaction in Pakistan [26]. These studies have assessed patient satisfaction from various angles, such as the provision of free medicine in public hospitals, comparing satisfaction between public and private hospitals, patient satisfaction at the medical, surgical, outdoor, registration departments, etc. For example, Farid *et al.*, studied the correlation between provision of free medicines to the patients in public hospitals and patient satisfaction in a cohort of 384 patients. The results showed that 59.4% of the respondents agreed that prices of medicines in Pakistan are low, 49.7% denied the availability of free medicines, 58.3% responded lack of proper information about drug utilization, while 63.3% respondents agreed that non-registration of pharmacies leads to substandard medicines [27]. Farooq *et al.*, compared patient satisfaction attending the Combined Military Hospital (CMH), Lahore and Jinnah Hospital, Lahore. The results revealed that patient satisfaction was significantly better ($p = 0.03$) at CMH in six (out of seven) domains studied [28]. Likewise, the patient satisfaction among private and public hospitals of Islamabad have also been compared in a multi-center cross-sectional study. The score of time spent by the doctor with the patient, communication, and convenience was significantly higher ($p < 0.05$) in private hospitals [29]. A recent study assessed patient satisfaction in tertiary care hospitals of Peshawar, with the use of PSQ-18. The results showed a higher patients satisfaction level for private hospitals compared to public hospitals (i.e., 96% vs. 86%) [30]. Similar comparison of patient satisfaction for public and private hospitals of Lahore has also been reported [31]. Moreover, patients' satisfaction with the utilization of different ambulance services at LRH Peshawar has been evaluated.

As a single-center cross-sectional study, the findings may have limited generalizability to other healthcare settings and only reflect patient perceptions at one point in time. The use of self-reported questionnaire data may introduce response bias and subjectivity. Future multicenter longitudinal studies with larger and more diverse populations are recommended to better explore causal relationships, incorporate qualitative feedback, and evaluate targeted interventions for improving technical quality, communication, and overall patient-centered care.

CONCLUSIONS

This study assessed the satisfaction of admitted patients and associated factors using PSQ-18 at the LRH, Peshawar, Pakistan. Overall, majority of the patients expressed satisfaction. Data stratification demonstrated that the highest number of patients were satisfied with the domain D7, followed by domain D6, while minimum number of patients were satisfied with the domain D2. Moreover, higher number of patients older than 45 years, patients of

female gender, patients with bachelor education and patient admitted to medical and allied ward expressed satisfaction. The same trend of patient satisfaction was found among male patients and female patients separately.

Authors' Contribution

Conceptualization: MW, KR, MP

Methodology: MP, JS, SMN

Formal analysis: MP, JS

Writing and Drafting: MW, KR, MP, JS, SMN

Review and Editing: MW, KR, MP, JS, SMN

All authors approved the final manuscript and take responsibility for the integrity of the work

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Alharbi HF, Alzahrani NS, Almarwani AM, Asiri SA, Alhwaymel FM. Patients' satisfaction with nursing care quality and associated factors: A cross-section study. *Nursing Open*. 2023 May; 10(5):3253-62. doi:10.1002/nop2.1577.
- [2] Moradi Rekabdar Kalaiee Z, Ghafouri R, Zandi M, Nasiri M. Effect of implementing of the IDEAL discharge model on satisfaction of patient referred to trauma emergency department. *PLOS One*. 2024 Jun; 19(6): e0304969. doi: 10.1371/journal.pone.0304969.
- [3] Maryam G, Sardar S, Khan J, Isram J, Hassan F. Factors affecting patient satisfaction during endoscopic procedures. *Scripta Medica*. 2024 Dec; 55(6): 749-54. doi: 10.5937/scriptamed55-51550.
- [4] Kanwel S, Ma Z, Li M, Hussain A, Erum N, Ahmad S. The influence of hospital services on patient satisfaction in OPDs: evidence from the transition to a digital system in South Punjab, Pakistan. *Health Research Policy and Systems*. 2024 Aug; 22(1):93. doi:10.1186/s12961-024-01178-8.
- [5] Huang H, Cui J, Zhang H, Gu Y, Ni H, Meng Y. The role of mental health in the relationship between nursing care satisfaction with nurse-patient relational care in Chinese emergency department nursing. *PLOS One*. 2024 Sep; 19(9): e0309800. doi:10.1371/journal.pone.0309800.
- [6] Abidova A, da Silva PA, Moreira S. Predictors of patient satisfaction and the perceived quality of healthcare in an emergency department in Portugal. *Western Journal of Emergency Medicine*. 2020 Jan; 21(2):391. doi: 10.5811/westjem.2019.9.44667.
- [7] Szasz TS and Hollender MH. A contribution to the philosophy of medicine: the basic models of the

- doctor-patient relationship. *AMA Archives of Internal Medicine*.1956May;97(5):585-92.doi:10.1001/archinte.1956.00250230079008.
- [8] Al-Hammouri F, Hamdan KM, Haymour AM, Ibrahim MO, Malkawi A, Al-Hiary SS et al. Quality of nursing care: Predictors of patient satisfaction in a national sample. *Nursing Open*.2024Aug;11(8):e2252.doi:10.1002/nop2.2252.
- [9] Gavurova B, Dvorsky J, Popesko B. Patient satisfaction determinants of inpatient healthcare. *International Journal of Environmental Research and Public Health*. 2021Oct; 18(21): 11337. doi: 10.3390/ijerph182111337.
- [10] Calhoun LM, Maytan-Joneydi A, Nouhou AM, Benova L, Delvaux T, van den Akker T et al. "I got what I came for": a qualitative exploration into family planning client satisfaction in Dosso region, Niger. *Open Access Journal of Contraception*.2022Jul;95-110.doi:10.2147/OAJC.S361895.
- [11] Jonkisz A, Karniej P, Krasowska DM. Meeting Patient Expectations: Assessing Medical Service and Quality of Care Using the SERVQUAL Model in Dermatology Patients at a Single Center in Poland. *Medical Science Monitor: International Medical Journal of Experimental and Clinical Research*.2023Aug;29:e941007-1.doi: 10.12659/MSM.941007.
- [12] Teshnizi SH, Aghamolaei T, Kahnouji K, Teshnizi SM, Ghani J. Assessing quality of health services with the SERVQUAL model in Iran. A systematic review and meta-analysis.*International Journal for Quality in Health Care*.2018Mar;30(2):82-9.doi:10.1093/intqhc/mzx200.
- [13] Evans DK and Welander Tärneberg A. Health-care quality and information failure: Evidence from Nigeria. *Health Economics*.2018Mar;27(3):e90-3.doi:10.1002/hec.3611.
- [14] Simsekler MC, Alhashmi NH, Azar E, King N, Luqman RA, Al Mulla A. Exploring drivers of patient satisfaction using a random forest algorithm. *BioMed Central Medical Informatics and Decision Making*.2021May;21(1): 157. doi: 10.1186/s12911-021-01519-5.
- [15] Thayaparan AJ and Mahdi E. The Patient Satisfaction Questionnaire Short Form (PSQ-18) as an adaptable, reliable, and validated tool for use in various settings. *Medical Education Online*.2013Jan;18(1):21747.doi:10.3402/meo.v18i0.21747.
- [16] Sullivan GM and Artino Jr AR. Analyzing and interpreting data from Likert-type scales. *Journal of Graduate Medical Education*.2013Dec;5(4):541-2.doi:10.4300/JGME-5-4-18.
- [17] Qadri SS, Pathak R, Singh M, Ahluwalia SK, Saini S, Garg PK. An assessment of patient's satisfaction with services obtained from a tertiary care hospital in rural Haryana. *International Journal of Collaborative Research on Internal Medicine and Public Health*. 2012 Aug; 4(8): 1524.
- [18] Woo S and Choi M. Medical service quality, patient satisfaction and intent to revisit: Case study of public hub hospitals in the Republic of Korea. *PLOS One*.2021 Jun;16(6):e0252241.doi:10.1371/journal.pone.0252241.
- [19] Wang MC, Chan PH, Paxton EW, Bellows J, Koplan K, Rabrenovich V et al. Factors influencing patient satisfaction with care and surgical outcomes for total hip and knee replacement.*The Permanente Journal*. 2021Nov; 25: 21-043. doi: 10.7812/TPP/21.043.
- [20] Duc Thanh N, My Anh BT, Xiem CH, Quynh Anh P, Tien PH, Thi Phuong Thanh N et al. Patient satisfaction with healthcare service quality and its associated factors at one polyclinic in Hanoi, Vietnam. *International Journal of Public Health*.2022Nov;67:1605055.doi:10.3389/ijph.2022.1605055.
- [21] Seo HL, Eom HK, Seo YK, Choi MJ, Im SI, Shin SM et al. Relationship among Nurses' Communication Style, Nursing Service Satisfaction, Hospital Service Satisfaction Perceived by Patients' Family and Intent to Revisit the Hospital. *Journal of Korean Clinical Nursing Research*. 2016; 22(3): 368-78.
- [22] Aiken LH, Sloane DM, Ball J, Bruyneel L, Rafferty AM, Griffiths P. Patient satisfaction with hospital care and nurses in England: an observational study. *British Medical Journal Open*.2021Jan;8(1):e019189.doi: 10.1136/bmjopen-2017-019189.
- [23] Ferreira PL, Raposo V, Tavares AI. Primary health care patient satisfaction: Explanatory factors and geographic characteristics. *International Journal for Quality in Health Care*.2020Mar;32(2):93-8.doi:10.1093/intqhc/mzz134.
- [24] Oster A, Wiking E, Nilsson GH, Olsson CB. Patients' expectations of primary health care from both patients' and physicians' perspectives: a questionnaire study with a qualitative approach. *BioMed Central Primary Care*.2024Apr;25(1):128.doi:10.1186/s12875-024-02389-2.
- [25] Hwang J, Vu GT, Tran BX, Nguyen TH, Nguyen BV, Nguyen LH et al. Measuring satisfaction with health care services for Vietnamese patients with cardiovascular diseases. *PLOS One*.2020Jun;15(6):e0235333. doi: 10.1371/journal.pone.0235333.
- [26] Naseer M, Zahidie A, Shaikh BT. Determinants of patient's satisfaction with health care system in Pakistan: a critical review. *Pakistan Journal of Public Health*. 2012 Jun; 2(2): 52.
- [27] Farid N, Shah M, Ali A, Jan D. Analyzing Patient Satisfaction With Regards To Provision Of Free Medicines In Khyber Pakhtunkhwa Pakistan. *Journal of Medical Sciences*.2021Mar;29(01):42-7.doi:10.52764/jms.21.29.1.11.
- [28] Farooq A, Khaliq MA, Toor MA, Amjad A, Khalid W, Butt F et al. Assessment of patient satisfaction in a military

- and public hospital: a comparative study. *Cureus*. 2020 Aug; 12(8). doi: 10.7759/cureus.10174.
- [29] Jan MU, Hassan Z, Khan MS, Ullah R, Siraj A. Comparison of Patient Satisfaction Level and its Various Determinants in Public and Private Hospitals in Peshawar. *Dr. Sulaiman Al Habib Medical Journal*. 2020 Dec; 2(4): 167-73. doi: 10.2991/dsahmj.k.200903.001.
- [30] Al-Hussami M, Al-Momani M, Hammad S, Maharmeh M, Darawad M. Patients' perception of the quality of nursing care and related hospital services. *Health and Primary Care*. 2017; 1(2): 1-6. doi: 10.15761/HPC.1000110.
- [31] Javed SA and Ilyas F. Service quality and satisfaction in healthcare sector of Pakistan-the patients' expectations. *International Journal of Health Care Quality Assurance*. 2018 Jul; 31(6): 489-501. doi: 10.1108/IJHCQA-08-2016-0110.