



Original Article



Screening Student Behavior: Exploring the Impact of Daily Screen Time on Sleep Quality, Mental Distress, and Academic Performance in Students

Kumayl Abbas Meghji^{1*}, Marvi Talpur², Aliza Khan², Hadiqa Fatima², Urooba Memon² and Monisha²

¹Department of Physiology, Isra University, Hyderabad, Pakistan

²Isra University, Hyderabad, Pakistan

ARTICLE INFO

Keywords:

Screen Time, Mental Health, Sleep Quality, Depression, Academic Performance

How to Cite:

Meghji, K. A., Talpur, M., Khan, A., Fatima, H., Memon, U., & Monisha, . (2025). Screening Student Behavior: Exploring the Impact of Daily Screen Time on Sleep Quality, Mental Distress, and Academic Performance in Students: Screening Student Behavior. *Pakistan Journal of Health Sciences*, 6(3), 152-158. <https://doi.org/10.54393/pjhs.v6i3.2745>

***Corresponding Author:**

Kumayl Abbas Meghji
 Department of Physiology, Isra University,
 Hyderabad, Pakistan
kumayl.abbas@isra.edu.pk

Received date: 10th January, 2025

Acceptance date: 19th March, 2025

Published date: 31st March, 2025

ABSTRACT

The pervasive influence of technology, particularly screen usage, on daily routines and its implications for sleep quality, mental health, academic performance, and physical activity has become an increasing area of concern. **Objective:** To evaluate the daily screen time of students from different fields of study and to assess its association with demographic variables, sleep quality, psychological symptoms, physical activity, and academic performance. **Methods:** This study was performed at Isra University, Hyderabad from February 2024 to August 2024, with 152 participants. Data on demographics, sleep quality (PSQI), mental distress (DASS-42), academic performance, and screen time were collected after informed consent. Statistical analysis, conducted using SPSS version 25.0, determined correlations between daily screen time, PSQI, and DAS scores. **Results:** The participants had a mean age of 20.59 ± 2.16 years, with most residing in urban areas (87.5%) and being day scholars (74.34%). Average daily screen time was 5.85 ± 1.14 hours, significantly higher among those with GPAs below 2.5 and low physical activity levels ($p < 0.05$). Mean scores for depression, anxiety, stress, and PSQI were 13.34 ± 12.01 , 11.17 ± 9.41 , 15.32 ± 11.09 , and 6.45 ± 3.48 , respectively. Screen time positively correlated with depression ($r = 0.81$, $p < 0.01$), anxiety ($r = 0.78$, $p < 0.01$), stress ($r = 0.83$, $p < 0.01$), and PSQI ($r = 0.75$, $p < 0.01$). **Conclusion:** Increased screen time was linked to poorer sleep quality, elevated mental distress, and reduced academic performance, particularly in urban populations and individuals using screens for recreation.

INTRODUCTION

With the rise of online connectivity, mobile devices, and social media platforms, technology has become an integral part of daily life, serving both recreational and professional purposes. This shift has led to a significant increase in the number of individuals using computers and electronic devices, thereby contributing to higher screen time [1, 2]. Prolonged exposure to screens emitting blue light disrupts the body's circadian rhythm, leading to sleep disturbances such as insomnia and irregular sleep patterns, which hinder sleep initiation and overall sleep quality [3]. Previous studies have shown that students with higher smartphone usage, particularly at night, experience poorer sleep quality and heightened anxiety levels [4,5]. Sleep plays a critical role in physical health, cognitive function, and

academic performance. Inadequate sleep has been associated with cardiometabolic disorders such as obesity, diabetes, and hypertension, as well as psychological issues like stress, depression, and anxiety [6]. Sleep deprivation impairs attention, mental processing speed, logical reasoning, and productivity, ultimately leading to declining academic performance [6]. Like poor sleep, depression is more prevalent among younger individuals with prolonged digital content consumption, particularly on computers [1]. Research suggests that excessive screen use, combined with social media-driven validation pressures, increases vulnerability to depression. This reliance on virtual interactions reduces face-to-face socialization, often leading to social withdrawal and

hormonal imbalances in dopamine, serotonin, and endorphins—neurotransmitters critical for emotional well-being [7]. While digital technology has revolutionized education, communication, and information accessibility, prolonged screen exposure is associated with negative health consequences, including sleep deprivation, social behavior disruptions, and academic decline [8]. The global prevalence of mobile internet device usage has increased exponentially, with a thousand-fold rise in the past decade [9]. Studies indicate that daily screen time exceeding two hours is now common and may adversely impact both physical and psychological health [10]. Excessive screen time has been strongly linked to declining academic performance, largely due to its effects on cognitive function, concentration, and sleep patterns. Research suggests that late-night screen exposure disrupts sleep cycles and reduces memory consolidation, impairing focus, learning efficiency, and information retention [11]. Additionally, screen-based activities, particularly social media and entertainment content, contribute to academic procrastination, reducing the time available for studying and completing assignments [12]. Studies have also found a direct relationship between increased screen time and lower GPA, with students reporting difficulty concentrating, increased fatigue, and reduced classroom engagement [13]. Given these concerns, understanding the role of screen time in academic success is essential for developing strategies to balance digital engagement with educational achievement. Although many studies have explored the effects of screen time on sleep and mental health, research focusing on university students remains limited, particularly in developing countries, where digital habits and academic pressures may differ. Furthermore, prior research often examines mental health and sleep disturbances separately, without considering how these factors collectively impact academic performance. This study aims to bridge these gaps by providing a comprehensive analysis of the relationship between screen time, sleep quality, mental distress, and academic performance among university students.

Excessive screen time among university students has become a growing public health concern due to its potential effects on sleep quality, mental health, physical activity, and academic performance. Although previous studies have investigated the relationship between screen exposure and psychological or sleep-related problems, limited research has comprehensively examined these factors together among university students in developing countries such as Pakistan, where digital behaviors and academic pressures may differ. Furthermore, the impact of recreational versus academic screen usage on students' well-being remains underexplored. This study's objective was to evaluate the daily screen time of students across

various fields of study and to assess its association with demographic variables, sleep quality, mental health symptoms, and academic achievement.

METHODS

This cross-sectional research was performed from February 2024 to August 2024, conducted at Isra University, Hyderabad after being approved by the Isra University Ethical Review Board (ERB letter # IU/RR-10-IRC-24/N/2024/108). The sample size was estimated using the OpenEpi calculator, with a 95% confidence level ($\alpha = 0.05$) and 80% statistical power ($\beta = 0.20$). An anticipated moderate correlation ($r \approx 0.3$) between screen time and mental health outcomes was used, based on previous studies that reported similar effect sizes [14]. To mitigate the impact of dropouts and non-responses, the initial estimate of 85 participants was adjusted upward by 20%, yielding a final target of 102 participants. However, a total of 152 students from various academic disciplines were recruited using purposive sampling, ensuring sufficient power for subgroup analyses and correlation studies. The sample included students from medical and allied health sciences (MBBS: 30, BDS: 20, DPT: 15, Pharmacy: 12) and non-medical fields (Engineering: 25, MBA: 25, BBA: 25), to capture a broad spectrum of screen time habits and their impacts. Informed consent was obtained after explaining the study objectives. The inclusion criteria were students who consented to participate, while those with pre-existing mental health conditions, undergoing psychological treatment, or unwilling to participate were excluded. This study controlled for certain confounding variables through eligibility criteria and participant selection. Students with pre-existing mental health conditions or those undergoing psychological treatment were excluded to minimize health-related confounders. Additionally, socioeconomic status was considered in the questionnaire, ensuring that only participants from similar educational and financial backgrounds were included, thereby reducing potential disparities. While other lifestyle factors were not explicitly controlled, the homogeneity of the sample helped minimize their impact. Demographic data included age, gender, residence, and campus accommodation. Academic performance data were self-reported by participants, providing their GPA as an indicator of their scholastic achievements. Participants provided consent for accessing their most recent GPA, which was verified through official academic records maintained by the institution. Physical activity levels were evaluated by asking participants about their frequency of exercise, categorized as frequent (≥ 3 times/week) or rare/none. Sleep quality was assessed using the Pittsburgh Sleep Quality Index (PSQI), a standardized self-reported

questionnaire that evaluates sleep disturbances over the past month. It consists of seven components, including subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleep medication, and daytime dysfunction. The total PSQI score ranges from 0 to 21, with higher scores indicating poorer sleep quality. The PSQI has demonstrated high internal consistency (Cronbach's $\alpha = 0.68$) and strong validity in measuring sleep disturbances [15]. Mental distress, including depression, anxiety, and stress, was assessed using the Depression, Anxiety, and Stress Scale (DASS-42), a widely validated self-reported instrument. It consists of 42 items divided into three subscales, each measuring symptoms of depression, anxiety, and stress over the past week. Each item is rated on a 4-point Likert scale (0 = Did not apply to me at all, 3 = Applied to me very much or most of the time), with higher scores indicating greater psychological distress. The total score is obtained by summing the responses for each subscale, with established cutoff points to categorize symptom severity. The DASS-42 has demonstrated excellent reliability, with Cronbach's $\alpha = 0.96$ [16]. Daily screen time data were self-reported, measured in hours per day, and categorized into <5 and ≥ 5 hours. Pearson's correlation coefficient was used to assess the linear relationship between the quantitative variables: daily screen time, PSQI, and DASS scores. The Chi-square test was applied to evaluate associations between the qualitative variables: screen time, demographic factors (gender, residence, accommodation type), GPA categories, and physical activity levels. Data were analyzed by SPSS version 25.0.

RESULTS

The mean age of the participants was 20.59 ± 2.16 years. The majority of the participants i.e. 50% belonged to the 18-20 years' age group while only 4.6% of participants belonged to the <18 -year age group. A vast majority of the participants belonged to urban areas (87.5%) and were day scholars (74.34%). Table 1 provides the demographic details of the study participants.

Table 1: Demographic variables of the study population (n=152)

Category	Frequency (%)
Age Group (Years)	
<18	07(4.60%)
18 - 20	76(50.00%)
21 - 22	47(30.92%)
>22	22(14.47%)
Gender	
Male	73(48.02)
Female	79(51.97)
Residence	
Urban	133(87.5)

Rural	19(12.5)
Accommodation	
Day Scholar	113(74.34)
Hostelite	39(25.65)
Field of Study	
Medical	77(50.65)
Non-medical	75(49.34)

The study participant's mean daily screen time was identified as 5.85 ± 1.14 hours. A statistically significant association was observed between daily screen time and participants from the urban population ($p < 0.05$), as shown in Table 2. Additionally, screen usage purpose was significantly associated with daily screen time, with participants engaging in recreational screen time being more prone to report higher usage compared to those using screens mainly for academic purposes ($p < 0.05$). GPA also showed a significant relationship, where students with a GPA below 2.5 exhibited significantly higher screen time compared to their peers with a GPA above 2.5 ($p < 0.05$). Similarly, exercise frequency was associated with screen time, as students who engaged in regular physical activity (≥ 3 times/week) reported lower screen time than those with rare or no physical activity ($p < 0.05$). However, no significant association was noted between daily screen time and gender or campus accommodation ($p > 0.05$).

Table 2: Association of Demographic Variables With Daily Screen Time

Variables	Category	Daily Screen time in hours		χ^2	p-Value
		<5	≥ 5		
Gender	Male	19	54	0.21	0.640
	Female	18	61		
Residence	Urban	28	105	6.25	0.010*
	Rural	9	10		
Accommodation	Day Scholar	29	84	0.41	0.510
	Hostelite	8	31		
GPA	>2.5	31	14	68.87	0.001*
	<2.5	6	101		
Screen Usage Purpose	Academic	26	18	40.60	0.001*
	Recreational	11	97		
Exercise Frequency	≥ 3 times/week	33	25	53.96	0.001*
	Rare/None	4	90		

The average PSQI score among participants was 6.45 ± 3.48 . A significant positive correlation ($r = 0.75$, $p < 0.010$) indicated that increased daily screen time was linked to poorer sleep quality.




Figure 1: Correlation of Daily Screen Time with Sleep Quality (PSQI Score)

The mean scored of depression, anxiety, and stress were found to be 13.34 ± 12.01 , 11.17 ± 9.41 , and 15.32 ± 11.09 respectively. Figure 2 shows the correlation of these mental health variables with daily screen time. There was a statistically significant positive relationship observed between screen time and depression ($r = 0.81$, $p < 0.010$), anxiety ($r = 0.78$, $p < 0.010$), and stress ($r = 0.83$, $p < 0.010$).




Figure 2: Correlation of Daily Screen Time with A) Depression B) Stress C) Anxiety

DISCUSSION

This study's objective was to evaluate the daily screen time of students across various fields of study and to assess its association with demographic variables, sleep quality, mental health symptoms, and academic achievement. These findings revealed a statistically significant association between daily screen time and urban residency. This is consistent with the explanation provided by Nedjar-Guerre *et al.*, who noted that urban families generally own more screen devices compared to their rural counterparts, contributing to higher screen time in urban populations [17]. However, this contrasts with Varadarajan *et al.*, and Wang *et al.*, who found no association between screen time and residential settings [18, 19]. These discrepancies may stem from differences in study populations, as this research focused on university students, whereas previous studies included younger age groups or mixed demographics. Additionally, variations in internet accessibility, urban infrastructure, and cultural

screen use habits across different regions may contribute to these conflicting findings. This study found no significant correlation between daily screen time and gender. However, research by Whiting *et al.*, reported higher screen time among boys compared to girls [20]. These differences could be influenced by cultural and social norms, where in some regions, males may engage more in gaming or entertainment-based screen use, whereas in others, screen exposure may be more evenly distributed due to educational demands or technological accessibility among both genders. The study identified a significant positive correlation between screen time and PSQI scores, implying that greater screen time correlates with lower sleep quality. This aligns with Maurya *et al.*, who demonstrated that excessive screen time among young adults raised the likelihood of sleep problems [21]. Similarly, Adamczewska-Chmiel *et al.*, found that individuals with problematic smartphone usage experienced shorter sleep durations and more frequent sleep deprivation due to decreased serum melatonin levels caused by smartphone-induced magnetic fields [22]. Moreover, prolonged screen exposure, especially before bedtime, delays sleep onset due to blue light exposure and impairs body regeneration processes [23, 24]. Guerrero *et al.*, suggested that excessive screen behaviors replace essential activities such as sleep, further emphasizing the importance of balanced screen time [23]. This research found a significant positive association between screen time and mental health parameters, including depression, anxiety, and stress, aligning with findings by Li *et al.*, and Adamczewska-Chmiel *et al.*, who reported that high screen time correlates with heightened anxiety and depressive symptoms [5, 22]. These effects can be explained by several neurophysiological mechanisms, including dopamine dysregulation, cortisol elevation, and melatonin suppression [7, 9, 25]. Digital media, particularly social media and gaming, overstimulates the dopaminergic reward system, leading to reduced dopamine receptor sensitivity, which contributes to mood instability, impulsivity, and depressive symptoms [26]. Additionally, prolonged screen exposure triggers chronic low-grade stress responses, causing sustained cortisol elevation, which is linked to increased anxiety, emotional dysregulation, and cognitive impairments [7]. Furthermore, blue light from screens suppresses melatonin production, disrupting circadian rhythms, reducing deep sleep, and increasing emotional reactivity, further compounding mental health challenges [27]. Boers *et al.*, demonstrated that each additional hour spent on online platforms significantly increases depressive symptoms among youth [28], while Ma *et al.*, highlighted the role of social media and television exposure in exacerbating depressive tendencies [29]. The present study highlighted that students who used screens primarily for recreational purposes were more likely to exhibit higher

screen times, poorer sleep quality, and elevated mental distress compared to those who used screens for academic purposes. Furthermore, a significant relationship was observed between lower GPA scores and increased screen time, suggesting that excessive screen exposure detracts from academic performance. This aligns with the results of Sapci *et al.*, who reported that higher screen time significantly reduces GPA [30]. Similarly, an association was noted between higher screen time and lower physical activity, suggesting a more sedentary lifestyle, which contrasts with the findings of Hadianfard *et al.*, who found that recreational screen activities (≥ 2 hours/day) were linked to increased physical activity [31]. The relationship between screen time and physical activity varies across studies due to differences in screen type, usage purpose, and study context. Television and gaming are linked to prolonged sedentary behavior, whereas smartphones may allow mobility (e.g., listening to music while exercising). Additionally, academic screen use may reduce activity, while fitness apps and social media challenges encourage it. These variations suggest that screen time's impact on physical activity is not universal and depends on multiple contextual factors. In light of these findings, it is evident that excessive screen time poses considerable risks to physical, mental, and academic well-being.

This study has several limitations, including its cross-sectional design, which limits the ability to establish causal relationships between screen time and health outcomes. The use of self-reported questionnaires may also introduce recall and reporting bias, particularly regarding screen time, sleep quality, and mental health symptoms. In addition, the study was conducted at a single institution with a relatively small sample size, limiting the generalizability of the findings. Future research should further explore these contextual factors to better understand the complexities of screen exposure and its effects. Although the study contributes to the field, its limitations must be acknowledged, particularly the small sample size, which may restrict the extent to which the findings can be generalized. Future investigations should aim to bridge these gaps to offer an in-depth understanding of the complex relationships between screen time, health, and academic performance.

CONCLUSIONS

The study concluded that increased screen time is strongly correlated with diminished sleep quality, amplified mental distress manifested as raised levels of depression, anxiety, and stress and lower academic performance. These effects were particularly pronounced among urban populations and students using screens predominantly for recreational purposes. These observations stress the significance of balancing screen usage to safeguard both physical and mental health while supporting academic success.

Authors' Contribution

Conceptualization: KAM, MT, AK, HF, UM, M

Methodology: MT, AK, HF, UM, M

Formal analysis: KAM

Writing and Drafting: KAM, MT, AK, HF, UM, M

Review and Editing: KAM, MT, AK, HF, UM, M

All authors approved the final manuscript and take responsibility for the integrity of the work

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Wang X, Li Y, Fan H. The associations between screen time-based sedentary behavior and depression: a systematic review and meta-analysis. *BioMed Central Public Health*. 2019 Dec; 19:1-9. doi:10.1186/s12889-019-7904-9.
- [2] Uqaili A, Qaiser F, Abbas K, Rahman AA. Smartphone use among young doctors and their impact on patients of Liaquat University Hospital Jamshoro. *Imp. Journal of Interdisciplinary Research*. 2017 Jan; 3: 161-4.
- [3] Elsheikh AA, Elsharkawy SA, Ahmed DS. Impact of smartphone use at bedtime on sleep quality and academic activities among medical students at Al-Azhar University at Cairo. *Journal of Public Health*. 2024 Nov; 32(11):2091-100. doi:10.1007/s10389-023-01964-8.
- [4] Ejaz W, Altay S, Naeem G. Smartphone use and well-being in Pakistan: Comparing the effect of self-reported and actual smartphone use. *Digital Health*. 2023 Jul; 9:20552076231186075. doi:10.1177/20552076231186075.
- [5] Li L, Zhang Q, Zhu L, Zeng G, Huang H, Zhuge J *et al.* Screen time and depression risk: A meta-analysis of cohort studies. *Frontiers in Psychiatry*. 2022 Dec; 13:1058572. doi: 10.3389/fpsy.2022.1058572.
- [6] Dos Santos AB, Prado WL, Tebar WR, Ingles J, Ferrari G, Morelhão PK *et al.* Screen time is negatively associated with sleep quality and duration only in insufficiently active adolescents: A Brazilian cross-sectional school-based study. *Preventive Medicine Reports*. 2024 Jan; 37:102579. doi:10.1016/j.pmedr.2023.102579.
- [7] Nakshine VS, Thute P, Khatib MN, Sarkar B. Increased screen time as a cause of declining physical, psychological health, and sleep patterns: a literary review. *Cureus*. 2022 Oct; 14(10). doi:10.7759/cureus.30051.

- [8] Hosen MJ, Eva SA, Rahman MM, Ibrahim M, Lira UF, Hossain AB et al. Health impacts of excessive use of Facebook among university students in Bangladesh. *Heliyon*.2021Jun;7(6).doi:10.1016/j.heliyon.2021.e07271.
- [9] Santos RM, Mendes CG, Sen Bressani GY, de Alcantara Ventura S, de Almeida Nogueira YJ, de Miranda DM et al. The associations between screen time and mental health in adolescents: a systematic review. *BioMed Central Psychology*.2023Dec;11(1):1-21.doi:10.1186/s40359-023-01166-7.
- [10] Lissak G. Adverse physiological and psychological effects of screen time on children and adolescents: Literature review and case study.*Environmental Research*.2018Jul;164:149-57.doi:10.1016/j.envres.2018.01.015.
- [11] Islam MR. Unplugging the threat: how internet addiction among adolescents undermines learning behavior. *SN Social Sciences*.2024Aug;4(9):161.doi:10.1007/s43545-024-00966-x.
- [12] Aslam I and Malik JA. Fear of Missing Out and Procrastination Among University Students: Investigating the Mediating Role of Screen Time. *CARC Research in Social Sciences*.2024Jun;3(2):206-13. doi:10.58329/criss.v3i2.122.
- [13] Gull M and Sravani BR. Do screen time and social media use affect sleep patterns, psychological health, and academic performance among adolescents? Evidence from bibliometric analysis.*Children and Youth Services Review*.2024Sep;164:107886.doi:10.1016/j.childyouth.2024.107886.
- [14] Lemola S, Perkinson-Gloor N, Brand S, Dewald-Kaufmann JF, Grob A. Adolescents' electronic media use at night, sleep disturbance, and depressive symptoms in the smartphone age.*Journal of Youth and Adolescence*.2015Feb;44(2):405-18.doi:10.1007/s10964-014-0176-x.
- [15] Zhang C, Zhang H, Zhao M, Li Z, Cook CE, Buysse DJ et al. Reliability, validity, and factor structure of Pittsburgh sleep quality index in community-based centenarians. *Frontiers in Psychiatry*.2020Aug;11:573530.doi:10.3389/fpsy.2020.573530.
- [16] Makara-Studzińska M, Tyburski E, Załuski M, Adamczyk K, Mesterhazy J, Mesterhazy A. Confirmatory factor analysis of three versions of the depression anxiety stress scale (DASS-42, DASS-21, and DASS-12) in Polish adults. *Frontiers in Psychiatry*.2022Jan;12:770532. doi:10.3389/fpsy.2021.770532.Nedjar-Guerre A,
- [17] Wattelez G, Serra-Mallol C, Frayon S, Galy O. Adolescent screen time and unhealthy food consumption in the context of the digital development in New Caledonia. *PLoS One*.2023May;18(5):e0285374. doi:10.1371/journal.pone.0285374.
- [18] Varadarajan S, Govindarajan Venguidesvarane A, Ramaswamy KN, Rajamohan M, Krupa M, Winfred Christadoss SB. Prevalence of excessive screen time and its association with developmental delay in children aged < 5 years: A population-based cross-sectional study in India.*PLoS One*.2021Jul;16(7):e0254102. doi:10.1371/journal.pone.0254102.
- [19] Wang H, Zhong J, Hu R, Fiona B, Yu M, Du H. Prevalence of high screen time and associated factors among students: a cross-sectional study in Zhejiang, China. *British Medical Journal Open*.2018Jun;8(6):e021493. doi:10.1136/bmjopen-2018-021493.
- [20] Whiting S, Buoncristiano M, Gelius P, Abu-Omar K, Pattison M, Hyska J et al. Physical activity, screen time, and sleep duration of children aged 6-9 years in 25 countries: An analysis within the WHO European childhood obesity surveillance initiative (COSI) 2015-2017.*Obesity Facts*.2021Dec;14(1):32-44.doi:10.1159/000511263.
- [21] Maurya C, Muhammad T, Maurya P, Dhillon P. The association of smartphone screen time with sleep problems among adolescents and young adults: cross-sectional findings from India.*BioMed Central Public Health*.2022Sep;22(1):1686.doi:10.1186/s12889-022-14076-x.
- [22] Adamczewska-Chmiel K, Dudzic K, Chmiela T, Gorzkowska A. Smartphones, the epidemic of the 21st century: a possible source of addictions and neuropsychiatric consequences. *International Journal of Environmental Research and Public Health*.2022 Apr;19(9):5152. doi:10.3390/ijerph19095152.
- [23] Guerrero MD, Barnes JD, Chaput JP, Tremblay MS. Screen time and problem behaviors in children: exploring the mediating role of sleep duration. *International Journal of Behavioral Nutrition and Physical Activity*.2019 Dec;16:1-0.doi:10.1186/s12966-019-0862-x.
- [24] Korcz A, Krzysztozek J, Bronikowski M, Łopatka M, Bojkowski Ł. Associations between physical activity, screen time, sleep time and selected academic skills in 8/9-year-old children. *BioMed Central Public Health*.2023 Jul;23(1):1335. doi:10.1186/s12889-023-16230-5.
- [25] Tang S, Werner-Seidler A, Torok M, Mackinnon AJ, Christensen H. The relationship between screen time and mental health in young people: A systematic review of longitudinal studies.*Clinical Psychology Review*.2021Jun;86:102021.doi:10.1016/j.cpr.2021.102021.
- [26] Verma A, Kumar A, Chauhan S, Sharma N, Kalani A, Gupta PC. Interconnections of screen time with neuroinflammation.*Molecular and Cellular Biochemistry*.2024 Sep:1-6.
- [27] Akman M. The Influence of Virtual Learning on the Rise in Screen Time Usage and Blue Light Exposure

- Ultimately Expanding Sleep Disturbances amongst Students. 2021 Apr.
- [28] Boers E, Afzali MH, Newton N, Conrod P. Association of screen time and depression in adolescence. *Journal of the American Medical Association Pediatrics*. 2019 Sep;173(9):853-9. doi:10.1001/jamapediatrics.2019.1759.
- [29] Ma L, Evans B, Kleppang AL, Hagquist C. The association between screen time and reported depressive symptoms among adolescents in Sweden. *Family Practice*. 2021 Dec;38(6):773-9. doi:10.1093/fampra/cmab029.
- [30] Sapci O, Elhai JD, Amialchuk A, Montag C. The relationship between smartphone use and students academic performance. *Learning and Individual Differences*. 2021 Jul;89:102035. doi:10.1016/j.lindif.2021.102035.
- [31] Hadianfard AM, Mozaffari-Khosravi H, Karandish M, Azhdari M. Physical activity and sedentary behaviors (screen time and homework) among overweight or obese adolescents: a cross-sectional observational study in Yazd, Iran. *BioMed Central Pediatrics*. 2021 Dec;21:1-0. doi:10.1186/s12887-021-02892-w.