



Original Article



Frequency of Obstetrical Anal Sphincter Injuries in Nulliparous Women Undergoing Normal Vaginal Delivery in Tertiary Care Hospital at Khyber Pakhtunkhwa

Nazish Farooq¹, Paghunda Roghani¹, Nuzhat Afza¹, Aisha Saleem¹ and Asma Farooq^{1*}

¹Department of Gynaecology and Obstetrics, Lady Reading Hospital, Peshawar, Pakistan

ARTICLE INFO

Keywords:

Obstetric Anal Sphincter Injury, Normal Vaginal Delivery, Nulliparous Women, Complications

How to Cite:

Farooq, N., Roghani, P., Afza, N., Saleem, A., & Farooq, A. (2025). Frequency of Obstetrical Anal Sphincter Injuries in Nulliparous Women Undergoing Normal Vaginal Delivery in Tertiary Care Hospital at Khyber Pakhtunkhwa: OASI in Nulliparous Women Undergoing Normal Vaginal Delivery. *Pakistan Journal of Health Sciences*, 6(3), 246-250. <https://doi.org/10.54393/pjhs.v6i3.2756>

***Corresponding Author:**

AsmaFarooq
 Department of Gynaecology and Obstetrics, Lady Reading Hospital, Peshawar, Pakistan
drasmafarooq10@gmail.com

Received date: 10th January, 2025

Acceptance date: 21st March, 2025

Published date: 31st March, 2025

ABSTRACT

Obstetric Anal Sphincter injury is one of the most devastating complications of vaginal delivery. The incidence is higher in instrumental deliveries, but OASI can be avoided with proper techniques, such as mediolateral episiotomy, and pre-assessment of fetal head position, such as vertex mal-presentation and progress of labour. **Objectives:** To find the frequency of Obstetric anal sphincter injury during normal vaginal deliveries in nulliparous female from August 2023 to September 2024. **Methods:** It was a cross-sectional study conducted in the Gynaecology and Obstetrics Department, Lady Reading Hospital. The study included 162 nulliparous women with Cephalic presentation undergoing Normal Vaginal Delivery, among them the patient with Obstetric Anal Sphincter Injury were identified to find the frequency. The data were collected by a structured questionnaire and analyzed with SPSS version 26.0. **Results:** Out of the total 162 populations, 6 (3.92%) individual had their OASIs. Five individuals had 3rd while one individual had a 4th-degree injury. The mean age was 26.33 years with a standard deviation of 4.73. The oldest mother was 38 years old, and the youngest was 18 years old. BMI from 18.5 to 21.9 (Kg/m²) was more frequent in a population of 67. Regarding the presentation of the fetus's head at birth, 146 presented Cephalic, 5 as transverse and 11 as oblique on presentation. **Conclusions:** It was concluded that the study shows the frequency of Obstetric Anal Sphincter injuries as 6 out of 162 individuals, having injuries of different degrees.

INTRODUCTION

Obstetric anal sphincter injuries (OASIS) occur after vaginal delivery. The internal and external anal sphincter and the anal mucosa are affected by these injuries, in more severe cases, which are referred to as third- and fourth-degree perineal lacerations [1]. OASIs are multifactorial among adolescents; it is dependent on maternal as well as fetal conditions. Diabetic mothers were considered at high risk [2]. In mothers of late age at first birth, increased blood pressure and birth weight of the baby were considered independent risk factors [3]. The risk of OASI can be evaluated before delivery by checking the mother's height, performing an ultrasound of the fetus' weight, determining the position of the occiput, and evaluating episiotomy

performance [4]. Two independent risk factors in women who have undergone previous vaginal delivery are only one previous vaginal delivery and the weight of the baby. The other risk factors were episiotomy and vacuum-assisted vaginal delivery [5-7]. Fecal incontinence is one of the consequences of OASI. The obstetric anal sphincter injury can be diagnosed with an ultrasound or a patient experiencing anal incontinence [8, 9]. To avoid injuring the obstetrical anal sphincter, it is recommended to select a delivery style that reduces the amount of trauma that occurs during the birth process. For instance, select a vacuum delivery rather than a forceps delivery, and take into consideration the mediolateral episiotomy technique



[10, 11]. Obstetrical anal sphincter injury can be prevented by identifying risk factors. Both antenatal and intrapartum obstetrical practice can lessen the injury, as prenatal massage from the last stage of pregnancy can improve the elasticity of muscles and give the perineum space to stretch [12]. Repair of third- and fourth-degree lacerations should follow a systematic approach, progressing from deep to superficial structures: the anorectal mucosa, anal sphincter complex, rectovaginal fascia, perineal body musculature, perineal skin, and vaginal muscular and epithelium [13]. A single continuous stitch is the most effective method of treatment for lacerations of the second degree. Lacerations of the anal sphincter complex require additional expertise, exposure, and lighting; it is recommended that attempts be made to relocate the procedure to the operating room [14]. This study was conducted to cover a population gap, as such a study had not been conducted in our demographics, and to see the frequency of OASIs in our ethnic group. In addition to this, our study has focused particularly on nulliparous women. Normal vaginal delivery was the focus of the current study. Although obstetric anal sphincter injuries (OASIS) are well studied globally, there is limited locally generated evidence from tertiary care settings in Khyber Pakhtunkhwa, particularly focusing on strictly selected nulliparous women undergoing spontaneous vaginal delivery. Most existing studies emphasize mixed obstetric populations or instrumental deliveries, leaving a gap in understanding the true baseline frequency of OASI in low-risk, primigravida women. Additionally, regional variations in obstetric practices, perineal protection techniques, and fetal presentation are not well documented in this population. The problem is that OASI remains an under-recognized but preventable complication of vaginal delivery, leading to long-term morbidity such as fecal incontinence and reduced quality of life. This study aims to determine the frequency of OASIS in women who are undergoing a normal vaginal delivery. By taking into consideration the risk factors that have been demonstrated in earlier research, this study also assists in determining which pregnant women are at risk for OASIS.

METHODS

A cross-sectional study was carried out including 162 individuals to assess the frequency of Obstetrical Anal Sphincter Injuries in nulliparous women, in the Department of Gynaecology and Obstetrics, Lady Reading Hospital-MTI, Peshawar. The study was conducted from August 2023 to September 2024. A non-probability consecutive sampling technique was used. The Ethical Approval of this research was obtained from IRB Lady Reading Hospital (LRH) Medical Teaching Institute (MTI), and the reference number was 727/LRH/MTI. The inclusion criteria used include All pregnant women having single intrauterine pregnancy,

normal vaginal delivery without instrumentation, only Primary-Gravida, no Previous History of 3rd or 4th Degree Tear, no Previous recto vaginal fistula repair, and patients delivered in Hospital Setting. The Exclusion criteria include the Fetus having a breech presentation, and women not willing to give consent were excluded from the study. Using the WHO sample size calculator, the sample size of the study was found, keeping the confidence interval as 95%, 7% absolute precision and a previously reported frequency of 29.2%. The sample size was 162 patients. After obtaining approval from the ethical committee of the hospital, the patients who were fulfilling the selection criteria, in the Gynaecology and Obstetrics Outpatient Department (OPD), LRH Peshawar. A Consultant Gynaecologist interviewed the patients along with the researcher. The aim of the study was discussed and explained to the patients, and then informed consent was taken. Demographics such as age, weight, height, period of gestation, and parity were noted. A detailed medical history was obtained, and a meticulous physical examination was performed. Relevant baseline labs, including CBC, RFTs, LFTs, ELECTROLYTES, FBS, VIRAL PROFILE, and U/S ABDOMEN, were sent to rule out any other pathology or multiple pregnancies. The researcher collected data and entered it into questionnaires. Confidentiality was ensured by masking the names of patients, and it was used for research purposes. Data entry and analysis were carried out using SPSS version 26.0. For quantitative variables like age, BMI, POG, fetal position during labor, and parity mean and standard deviation were calculated. Frequency and percentage were calculated for categorical variables like gender and type of previous delivery. The BMI, gestational age at delivery, and fetal presentation at birth were stratified and shown in tables. Finally, the mode of induction of labour was measured as frequency.

RESULTS

The mean age of the population was 26.33 with an SD of 4.73. The youngest among the mothers was 18 years old, while the oldest was 38 years old. The weight and height of mothers are used in measuring BMI (Kg / m^2) among them 21 mothers have BMI less the 18.5 (Kg / m^2), 67 mothers have BMI from 18.5 to 21.9 (Kg / m^2), 59 have a BMI range from 22 to 24.9 (Kg / m^2), while 12 mothers have BMI of 25 to 30 (Kg / m^2). And only one mother has a BMI of more than 30 (Kg / m^2) (Table 1).

Table 1: Demographics Presentation of participants(n=162)

Variables	Mean \pm SD/ n (%)
Age (Years)	26.33 + 4.73
Minimum	18.00
Maximum	38.00
BMI (Kg / m^2)	
Less the 18.5	21 (12.96%)

18.5 to 21.9	67(43.35%)
22 to 24.9	59(36.41%)
25 to 29.9	12(7.4%)
More Than 30	1(0.61%)

Only term mothers were included in the study, which was divided into two groups: those with gestational age at delivery from 37 to 42 weeks, which had 155 individuals, and those with gestational age at delivery more than 42 weeks, which had 7 individuals. The presentation of the fetus's head at birth is shown in Figure 3: 146 presented cephalic, 5 presented transverse, and 11 presented oblique (Table 2).

Table 2: Gestational Age at Delivery and Fetal Presentation at Birth

Variables		Number of Mothers
Gestational Age at Delivery	37 to 42 weeks	155
	More than 42 weeks	7
Variables		
Fetal Presentation at birth	Cephalic	146
	Transverse	5
	Oblique	11

Out of the total 162, population, 6 (3.92%) individual had their OASIs. 5 individuals had 3rd degree and only one had a 4th-degree injury. So the total frequency of OASI in our study population was 6 out of 162 primigravida mothers (Table 3).

Table 3: Frequency of OASIS

Frequency of OASIS		
Total Number of NVDs (162)	Yes	No
3rd Degree	5	—
4th Degree	1	—
Total	6(3.92%)	156

DISCUSSION

Our study was conducted on a very specific population i.e. primigravida, full-term having a normal vaginal delivery. It was conducted in a closed, monitored environment. The frequency of total OASIs is 9 out of 162 primigravida mothers. A study by Sørbye et al., shows that women from South Asia had the highest incidence of OASI at 6.2%, followed by those from Southeast Asia, East Asia, and the Pacific at 5.7%, and Sub-Saharan Africa at 5.2% [15]. Kwok and fellows studied the prevalence of OASI in nulliparous women as 7.8% undergoing normal vaginal delivery [16]. According to a study by Marschalek, migrant women encounter a novel physical and social environment, where limited language proficiency significantly hinders social integration, leading to inadequate health literacy and suboptimal care post-migration [17]. Longo and his co-worker studied risk factors that prone the patient to OASIs undergoing normal vaginal delivery. They observed that the risk was twice for vacuum delivery in nulliparous women and the risk decreased with previous vaginal delivery and

also in spontaneous vaginal delivery. Epidural Anesthesia was considered preventive as it lowered the incidence of OASIS. Malpresentation of the fetal head and circumference of the fetal head were considered fetal risk factors [18]. Johannesson et al., studied the mixed response of women regarding the role of physiotherapy in recovery from obstetric anal sphincter injury some women felt difficulty and tiresome about therapy, while others were very happy with the outcome and find themselves lucky to have it [19]. A study shows that Anal sphincter rips are frequently overlooked after delivery, and even when they are identified, they are frequently not well healed, leaving a high percentage of residual abnormalities following reconstruction, according to mounting data. Between 25% and 50% of individuals will continue to experience anal incontinence even after initial repair and postpartum diagnosis [20]. Antonakou, studied that women suffering from anal incontinence after OASIs feels awkward and ashamed and hesitate to seek medical attention and the condition is termed as OASIs syndrome [21]. In a cohort study, Eggebø and fellows, showed that lateral episiotomy is associated with a lower incidence of OASIs in nulliparous women [22]. It is a study that mediolateral or lateral episiotomy is preventive for OASIs, decreasing the prevalence by 45%, while Forceps delivery is associated with a higher incidence of OASIs, the risk of OASIs is reduced to half by using vacume instead of forcep delivery (aOR = 1.92, 95% Cis = 1.79-2.05) [23]. The strength of our study was that it focused on single intrauterine primigravida females delivering babies via normal vaginal delivery in term babies. So, it helps us access the accurate frequency of this group and provides ground for further study.

This study is limited by its single-center design and relatively small sample size, which may reduce generalizability to other populations. The use of non-probability sampling may introduce selection bias, and important confounders such as duration of labor, episiotomy technique, and operator skill were not fully controlled. Additionally, long-term outcomes of OASI such as anal incontinence were not assessed. Future studies should include multicenter designs with larger sample sizes and more robust adjustment for confounding variables. Prospective cohort studies are recommended to better evaluate causal risk factors and outcomes. Furthermore, research focusing on preventive obstetric interventions such as episiotomy techniques, perineal support methods, and intrapartum risk scoring systems should be prioritized.

CONCLUSIONS

It was concluded that the frequency of obstetrical anal sphincter injuries in women undergoing normal vaginal delivery for nulliparous women in our study is 6 out of 162,

(3.92%). However, further research is required in the field with controlled confounding factors.

Authors' Contribution

Conceptualization: NF

Methodology: NF, NA, AS

Formal analysis: NF, PR, AF

Writing and Drafting: AF

Review and Editing: AF, NF, NA, AS

All authors approved the final manuscript and take responsibility for the integrity of the work

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Jan D, Jaromír M, Kamil Š, Alois M. Obstetric anal Sphincter Injury and Anal Incontinence. *Ceska Gynekologie*. 2021Jan;86(2):118-22. doi:10.48095/cccg2021118.
- [2] Meyer R and Levin G. Risk Factors for Obstetric Anal Sphincter Injury Among Adolescents. *European Journal of Obstetrics and Gynaecology and Reproductive Biology*. 2022Mar;270:151-5. doi:10.1016/j.ejogrb.2022.01.009.
- [3] Walker TV, Bryson C, Rahman S, Carter-Brooks CM. Risk of Obstetric Anal Sphincter Injury by Delivering Provider. *Reproductive Medicine*. 2024May;5(2):57-64. doi:10.3390/reprodmed5020007.
- [4] Meyer R, Schwartz A, Horesh N, Alcalay M, Ram E, Levin G. A Dynamic Prediction Score for Obstetric Anal Sphincter Injury Among Nulliparous Women Delivering Vaginally. *International Journal of Gynaecology and Obstetrics*. 2022May;157(2):271-6. doi:10.1002/ijgo.13730.
- [5] Levin G, Rottenstreich A, Tsur A, Cahan T, Yoeli-Ullman R, Shai D et al. Risk Factors for Obstetric Anal Sphincter Injury Among Parous Women. *Archives of Gynaecology and Obstetrics*. 2021Mar;303:709-14. doi:10.1007/s00404-020-05806-w.
- [6] Chill HH, Karavani G, Lipschuetz M, Yishai K, Winer J, Shimonovitz T et al. Birthweight Difference Between Deliveries and the Risk Of Obstetric Anal Sphincter Injury In Parous Women. *International Urogynecology Journal*. 2022 Dec;33(12):3401-6. doi:10.1007/s00192-022-05207-4.
- [7] Rygaard A, Jonsson M, Wikström AK, Brismar-Wendel S, Hesselman S. Obstetric anal Sphincter Injury in the Second Birth After Perineal Wound Complication in the First Birth: A Nationwide Register Cohort Study. *An International Journal of Obstetrics and Gynaecology*. 2024 Sep;131(10):1378-84. doi:10.1111/1471-0528.17830.
- [8] Bharucha AE, Knowles CH, Mack I, Malcolm A, Oblizajek N, Rao S et al. Faecal Incontinence in Adults. *Nature Reviews Disease Primers*. 2022Aug;8(1):53. doi:10.1038/s41572-022-00381-7.
- [9] Sideris M, McCaughey T, Hanrahan JG, Arroyo-Manzano D, Zamora J, Jha S et al. Risk of Obstetric Anal Sphincter Injuries (OASIS) and Anal Incontinence: A Meta-Analysis. *European Journal of Obstetrics and Gynaecology and Reproductive Biology*. 2020 Sep;252:303-12. doi:10.1016/j.ejogrb.2020.06.048.
- [10] Gachon B, Becam E, Barussaud ML, Carlier-Guerin C, Fritel X. How Can We Improve Our Practices in Obstetric Anal Sphincter Injury Prevention, Diagnosis, and Management of Symptomatic Women? *Journal of Gynaecology, Obstetrics and Human Reproduction*. 2021Nov;50(9):102183. doi:10.1016/j.jogoh.2021.102183.
- [11] Luchrist D, Brown O, Pidaparti M, Kenton K, Lewicky-Gaupp C, Miller ES. Predicting Obstetrical Anal Sphincter Injuries in Patients Who Undergo Vaginal Birth After Cesarean Delivery. *American Journal of Obstetrics and Gynaecology*. 2021Aug;225(2):173-e1. doi:10.1016/j.ajog.2021.02.014.
- [12] Okeahialam NA, Sultan AH, Thakar R. The Prevention of Perineal Trauma During Vaginal Birth. *American Journal of Obstetrics and Gynaecology*. 2024Mar;230(3):S991-1004. doi:10.1016/j.ajog.2022.06.021.
- [13] Schmidt PC and Fenner DE. Repair of Episiotomy and Obstetrical Perineal Lacerations (First-Fourth). *American Journal of Obstetrics and Gynaecology*. 2024Mar;230(3):S1005-13. doi:10.1016/j.ajog.2022.07.005.
- [14] Arnold MJ, Sadler K, Leli K. Obstetric Lacerations: Prevention and Repair. *American Family Physician*. 2021Jun;103(12):745-52.
- [15] Sørbye IK, Bains S, Vangen S, Sundby J, Lindskog B, Owe KM. Obstetric anal Sphincter Injury by Maternal Origin and Length of Residence: A Nationwide Cohort Study. *An International Journal of Obstetrics and Gynaecology*. 2022Feb;129(3):423-31. doi:10.1111/1471-0528.16985.
- [16] Kwok SP, Wan OY, Cheung RY, Lee LL, Chung JP, Chan SS. Prevalence of Obstetric Anal Sphincter Injury Following Vaginal Delivery in Primiparous Women: A Retrospective Analysis. *Hong Kong Medical Journal*. 2019 Aug;25(4):271. doi:10.12809/hkmj197842.
- [17] Marschalek ML. Rates of Obstetric Anal Sphincter Injuries Among Immigrant Women. *An International Journal of Obstetrics and Gynaecology*. 2022Feb;

- 129(432):1471-0528.doi:10.1111/1471-0528.16991.
- [18] Longo VL, Odjidja EN, Zanfini BA, Catarci S, Carducci B, Draisci G *et al.* Risk Factors Associated with Severe Perineal Lacerations During Vaginal Delivery: A 10-Year Propensity Score-Matched Observational Study. *American Journal of Obstetrics and Gynaecology Global Reports*. 2023 May; 3(2):100174. doi:10.1016/j.xagr.2023.100174.
- [19] Johannesson E, Sjöberg ÅL, Segerbrand N, Olsén MF, Gutke A. Women's Experiences of Obstetric Anal Sphincter Injury and Physical Therapy Interventions- A Qualitative Study. *Brazilian Journal of Physical Therapy*. 2022 Mar; 26(2):100397. doi:10.1016/j.bjpt.2022.100397.
- [20] Bellussi F and Dietz HP. Postpartum Ultrasound for the Diagnosis of Obstetrical Anal Sphincter Injury. *American Journal of Obstetrics and Gynaecology Maternal-Fetal Medicine*. 2021 Nov; 3(6):100421. doi: 10.1016/j.ajogmf.2021.100421.
- [21] Antonakou A. The Long-Term Physical, Emotional and Psychosexual Outcomes Related to Anal Incontinence After Severe Perineal Trauma at Childbirth. *European Journal of Midwifery*. 2018 Aug; 2: 8. doi: 10.18332/ejm/93544.
- [22] Eggebø TM, Rygh AB, von Brandis P, Skjeldestad FE. Prevention of Obstetric Anal Sphincter Injuries with Perineal Support and Lateral Episiotomy: A Historical Cohort Study. *Acta Obstetrica et Gynaecologica Scandinavica*. 2024 Mar; 103(3):488-97. doi: 10.1111/aogs.14742.
- [23] Fodstad K, Laine K, Räisänen S. Obstetric anal Sphincter Injuries During Instrumental Vaginal Delivery: An Observational Study Based On 18-Years of Real-World Data. *An International Journal of Obstetrics and Gynaecology*. 2024 Dec; 131(13):1824-31. doi:10.1111/1471-0528.17914.